

Geometric motives and the h-topology

Jakob Scholbach

ABSTRACT

This paper reinterprets Voevodsky's category of geometric motives. Using the h-topology we obtain an equivalent description of the category avoiding correspondences.

Introduction

The category of geometric motives [Voe00b] is roughly defined to be the localisation of the category of smooth correspondences with respect to the affine line and the Zariski topology. Our principal result is an alternative approach to this category: with rational coefficients it is possible to avoid the correspondences by replacing the Zariski topology by the much finer h-topology, where the base field is supposed to be of characteristic zero.

We see two principal facets of this description: one the one hand it clarifies why the technique of proper descent, which is often applied by replacing some scheme by a proper hypercovering in order to resolve its singularities, fits into the motivic framework (e.g. in [GS96]). As a byproduct of this phenomena, the whole category of schemes naturally maps to the category of motives. On the other hand, the theorem is a hint that correspondences, which make the handling of the several categories of motives technically intricate, should not be necessary in the case of rational coefficients. This might help to study motives over more general bases.

We define a category of *effective quasigeometric motives* with rational coefficients as follows: Let \mathbf{Sch} denote the category of separated schemes of finite type over a field k . Let $\mathbb{Q}\mathbf{Sch}$ be the \mathbb{Q} -linear hull of \mathbf{Sch} (i.e. morphisms are given by \mathbb{Q} -linear combinations of morphisms of schemes) and $\overline{\mathbb{Q}\mathbf{Sch}}$ its closure under countable direct sums. Finally let $\mathbf{K}^- (\overline{\mathbb{Q}\mathbf{Sch}})$ denote the homotopy category of bounded above complexes. We formally invert the following complexes:

- $X \times \mathbb{A}^1 \rightarrow X$, X smooth
- Čech nerves of h-coverings: $\dots \rightarrow U \times_X U \rightarrow U \rightarrow X$, for U smooth, $X \in \mathbf{Sch}$, $U \rightarrow X$ an h-covering.

The h-topology is essentially given by Zariski open coverings, finite surjective morphisms and blowups. We define the category of *effective quasigeometric motives* to be the localisation

$$\mathbf{DM}_{\text{gm}}^{\text{eff}}(\mathbb{Q}) := \mathbf{K}^- (\overline{\mathbb{Q}\mathbf{Sch}}) / \{ \mathbb{A}^1, \text{h-coverings} \}$$

The subcategory $\mathbf{DM}_{\text{gm}}^{\text{eff}}(\mathbb{Q})$ of bounded complexes of schemes is called category of *effective geometric motives*.

On the other hand, the category of effective geometrical motives with transfers is defined via the category $\mathbb{Q}\mathbf{SmCor}$, consisting of smooth schemes over k and morphisms given by finite correspondences, i.e. \mathbb{Q} -linear combinations of cycles $W \subset X \times Y$ finite over X . Again, we first define a quasigeometric category by

$$\mathbf{DM}_{\text{gm, tr}}^{\text{eff}}(\mathbb{Q}) := \mathbf{K}^- (\overline{\mathbb{Q}\mathbf{SmCor}}) / \{ \mathbb{A}^1, \text{Mayer-Vietoris} \},$$

where $\overline{\mathbf{QSmCor}}$ is the closure of \mathbf{QSmCor} under countable direct sums and Mayer-Vietoris complexes are given by $U \cap V \rightarrow U \sqcup V \rightarrow U \cup V$ (U, V smooth). Its subcategory $\mathbf{DM}_{\text{gm, tr}}^{\text{eff}}(\mathbb{Q})$ of complexes in $\mathbf{K}^b(\mathbf{QSmCor})$ is Voevodsky's category of *effective geometric motives with transfers*.

Our principal theorem is the following:

Theorem 4.1. *Let k be of characteristic zero. Then there is an equivalence of categories of effective geometric motives with and without transfers (with rational coefficients):*

$$\mathbf{DM}_{\text{gm, tr}}^{\text{eff}}(\mathbb{Q}) \xrightarrow{\cong} \mathbf{DM}_{\text{gm}}^{\text{eff}}(\mathbb{Q}).$$

This equivalence derives from an analogous equivalence of the quasigeometric variants. These equivalences extend to the non-effective motivic categories (see 4.4 for their definition).

This theorem is a considerable simplification of geometric motives as morphism groups are easier to calculate. Moreover, the category naturally contains all schemes and the h-topology permits greater flexibility.

Overview of the proof and of the paper: A principal notion are *presheaves with transfers*. These are just contravariant functors $\mathbf{SmCor} \rightarrow \mathbf{Ab}$. A presheaf is called homotopy invariant, if the canonical map $F(X) \rightarrow F(X \times \mathbb{A}^1)$ is an isomorphism.

The general idea is to refine the topology to avoid correspondences. The h-topology is well suited for this task: firstly, every h-sheaf has a unique transfer structure (Proposition 2.2). Secondly, for an homotopy invariant presheaf with transfers with rational coefficients, the sheafification with respect to the Zariski-, Nisnevich-, étale, cdh-, qfh- or h-topology agree. Moreover, its cohomologies with respect to these topologies agree (Theorem 2.11).

These facts rather directly give an equivalence of categories of *motivic complexes* with rational coefficients, with and without transfers, respectively (Theorem 3.7):

$$\mathbf{DM}_{\text{Nis, tr}}^{\text{eff}, -}(\mathbb{Q}) \cong \mathbf{DM}_{\text{h}}^{\text{eff}, -}(\mathbb{Q}),$$

where the former category is the full subcategory of the derived category of Nisnevich sheaves with transfers $\mathbf{D}^-(\mathbf{Shv}_{\text{Nis, tr}}(\mathbf{Sm}))$ consisting of complexes with homotopy invariant cohomology-Nisnevich-sheaves and the latter is defined similarly using h-sheaves instead.

Applying the cohomology comparison results, we obtain the embedding theorem (3.14):

$$\mathbf{DM}_{\text{gm}}^{\text{eff}}(\mathbb{Q}) \subset \mathbf{DM}_{\text{gm}}^{\text{eff}}(\mathbb{Q}) \subset \mathbf{DM}_{\text{h}}^{\text{eff}, -}(\mathbb{Q}).$$

The same statement is known for the situation with transfers (Theorem 3.15). More precisely, in either the situation with or without transfers, the geometric category is the subcategory of compact objects and the quasigeometric category is the one of \aleph_1 -compact objects (Definition A.3). This completes the proof.

We restrict to characteristic zero to be able to resolve singularities (see Proposition 1.1). We work with \mathbb{Q} -coefficients in order to ensure the finiteness of the cohomological dimension and the equality of Nisnevich and étale cohomologies. Another technically important point is Lemma 2.6, which is used to close the gap – essentially consisting of finite surjective morphisms – between Nisnevich and qfh-topology. Analogous cohomology comparison results for sheaves with integral coefficients are not true (see Remark 2.12).

The proof of the theorem heavily relies on results and techniques developed by Friedlander, Suslin and Voevodsky, especially the cohomology comparison results and sheaves with transfers [VSF00, Voe96, SV96]. Another technically important tool is Neeman's compactness notion [Nee01],

slightly adapted to our situation.

I was writing this paper under the supervision of Annette Huber [Sch05]. I would like to thank her for her constant (in)valuable support.

1. Notations and Preliminary Results

We fix the following notations. The base field is denoted by k . By scheme, we always mean a separated scheme of finite type over $\text{Spec } k$. We denote by $\mathbf{Sm} \subset \mathbf{Sch}$ the categories of (smooth) schemes and by $\mathbb{Z}\mathbf{Sch}$ the category whose objects are the same as in \mathbf{Sch} , but morphisms are given by $\text{Hom}_{\mathbb{Z}\mathbf{Sch}}(X, Y) := \bigoplus_i \mathbb{Z}[\text{Hom}_{\mathbf{Sch}}(X_i, Y)]$, where X_i are the connected components of X . By formally adding a zero object, we obtain an additive category.

Recall the definition of Voevodsky's category of finite correspondences \mathbf{SmCor} (see, for example [MVW, esp. Lecture 1]): its objects are smooth schemes. Its morphisms are defined by

$$\text{Hom}_{\mathbf{SmCor}}(X, Y) := \mathbb{Z}[W \subset X \times Y, W \text{ irreducible, surjective and finite over } X],$$

if X is connected and $\text{Hom}_{\mathbf{SmCor}}(\sqcup X_i, Y) := \bigoplus \text{Hom}_{\mathbf{SmCor}}(X_i, Y)$ otherwise. The composition of correspondences is defined in the usual way (see *loc. cit.*). We obtain an additive category. Assigning to a morphism in \mathbf{Sm} its graph gives a faithful functor $\mathbf{Sm} \rightarrow \mathbf{SmCor}$ (see [Ful98, §16] for associativity of the composition of correspondences and functoriality).

Presheaves are denoted by $\mathbf{PSh}(-)$, where $-$ is either \mathbf{Sm} or \mathbf{Sch} . If not mentioned explicitly otherwise, presheaves will be presheaves of abelian groups. All our presheaves will be understood to be additive, i.e. $F(X \sqcup Y) \cong F(X) \oplus F(Y)$. If t is a topology, t -sheaves are denoted by $\mathbf{Shv}_t(-)$. The t -sheafification of a presheaf F is denoted by F_t . In general we add a subscript $-\text{tr}$ when there is some category or object “with transfers”: the categories of t -sheaves with transfers (i.e. contravariant functors from \mathbf{SmCor} to abelian groups whose restriction to \mathbf{Sm} is a t -sheaf) are denoted by $\mathbf{Shv}_{t,\text{tr}}(\mathbf{Sm})$ and similarly for presheaves with transfers: $\mathbf{PSh}_{\text{tr}}(\mathbf{Sm})$.

The representable presheaves on schemes are written $\mathbb{Z}(-)$ (and $\mathbb{Z}_{\text{tr}}(-)$ for correspondences, i.e. $\mathbb{Z}_{\text{tr}}(X)(V) = \text{Hom}_{\mathbf{SmCor}}(V, X)$).

The \mathbb{Q} -rational variants of these constructions are denoted by $\mathbb{Q}\mathbf{Sch}$, $\mathbb{Q}\mathbf{SmCor}$, \mathbb{Q}_{tr} etc.

Let $\mathbf{Kom}^*(-)$, $*$ $\in \{-, b\}$ denote the category of bounded above and bounded complexes, respectively (with differentials having degree $+1$), $\mathbf{K}^*(-)$ the corresponding homotopy category and $\mathbf{D}^*(-)$ the derived category. The total complex of a double complex is denoted by $\text{Tot}(-)$ (see [Wei94, 1.2.6] for choice of signs). If \mathcal{C} , \mathcal{D} are categories, $\mathcal{C} \subset \mathcal{D}$ denotes a full embedding.

We say “ k admits weak resolution of singularities”, if every scheme X admits a proper surjective morphism $Y \rightarrow X$, where Y is smooth.

We say “ k admits strong resolution of singularities”, if furthermore for any abstract blow-up $p : Y \rightarrow X$ with (closed) center $Z \subset X$ of any scheme X (i.e. any proper surjective map, such that $p^{-1}(X - Z) \cong X - Z$) there is a finite composition of blow-ups of *smooth* centers $q : \tilde{X} = X_n \rightarrow \dots \rightarrow X_0 = X$ factoring through p .

PROPOSITION 1.1. *Any field admits weak resolution of singularities. Any field of characteristic 0 admits strong resolution of singularities.*

Proof. For the weak resolution, see [Hir64, Main Theorem I, S. 132] (char $k = 0$) and [dJ96, Theorem 4.1; §1] (char $k > 0$).

For the strong resolution over a field of characteristic zero we apply the resolution of indeterminacies to p^{-1} , see [Hir75] in the analytical setting and [AKMW02, §1.2.4], [RG71, Corollaire 5.7.12], [Bon02, §1.2] for the algebraic setting. \square

DEFINITION 1.2. For a simplicial scheme \mathcal{X} we denote by $\mathbb{Z}(\mathcal{X})$ the Moore complex of representable presheaves

$$\dots \longrightarrow \mathbb{Z}(\mathcal{X}_2) \xrightarrow{\delta_2} \mathbb{Z}(\mathcal{X}_1) \xrightarrow{\delta_1} \mathbb{Z}(\mathcal{X}_0),$$

where $\delta_n = \sum_{i=0}^{n+1} (-1)^i \partial_n^i$ and $\partial_n^i : \mathcal{X}_n \rightarrow \mathcal{X}_{n-1}$ is the i -th boundary map of \mathcal{X} .

For a morphism $U \rightarrow X$ we denote by \mathcal{U}_X the simplicial scheme $\dots \rightarrow U_X^3 \rightarrow U_X^2 \rightarrow U$ with the obvious boundary and degeneracy maps. If $U \rightarrow X$ is a covering with respect to a (pre-)topology t , \mathcal{U}_X is a t -hypercovering of X .

LEMMA 1.3. Let $\mathcal{X} = (X_n)_{n \in \mathbb{N}}$ be a hypercovering in $(\mathbf{Sch}/X)_t$, t an arbitrary topology. The augmented Moore complex of the corresponding representable t -sheaves $\mathbb{Z}_t(\mathcal{X})$ is exact:

$$\dots \rightarrow \mathbb{Z}_t(\mathcal{X}_2) \rightarrow \mathbb{Z}_t(\mathcal{X}_1) \rightarrow \mathbb{Z}_t(\mathcal{X}_0) \rightarrow \mathbb{Z}_t(X) \rightarrow 0.$$

Proof. A theorem of Verdier [SGA72, Exposé V, Théorème 7.3.2(3)] asserts that for any hypercovering \mathcal{X} in $(\mathbf{Sch}/X)_t$ the sheafification of the following complex is exact:

$$\dots \rightarrow \mathbb{Z}_X(\mathcal{X}_2) \rightarrow \mathbb{Z}_X(\mathcal{X}_1) \rightarrow \mathbb{Z}_X(\mathcal{X}_0) \rightarrow \mathbb{Z}_X(X) = \mathbb{Z} \rightarrow 0,$$

where $\mathbb{Z}_X(U)(V) := \mathbb{Z}[\mathrm{Hom}_{\mathbf{Sch}/X}(V, U)]$. The proof of *loc. cit.* is valid in our situation, if we replace $\mathbb{Z}_{X,t}(-)$ by $\mathbb{Z}_{\mathrm{Spec} k,t}(-) = \mathbb{Z}_t(-)$. \square

LEMMA 1.4. Let $f : U \rightarrow X$ be an arbitrary morphism. Then the complex

$$\mathbb{Z}(\mathcal{U}_X) \rightarrow \mathbb{Z}(X) \rightarrow \mathrm{coker} \mathbb{Z}(f) \rightarrow 0$$

is an exact complex (of presheaves).

Proof. This is [Voe96, Proposition 2.1.4]. We give a proof for the convenience of the reader. Let A be an arbitrary connected scheme. For brevity, $(A, -)$ and $[A, -]$ denote $\mathrm{Hom}_{\mathbf{Sch}}(A, -)$ and $\mathbb{Z}(-)(A) = \mathrm{Hom}_{\mathbb{Z}\mathbf{Sch}}(A, -)$, respectively. We have to show that the following complex is exact:

$$\dots \rightarrow [A, U_X^3] \rightarrow [A, U_X^2] \rightarrow [A, U] \rightarrow [A, X] \rightarrow \mathrm{coker} \mathbb{Z}(f)(A) \rightarrow 0.$$

We may suppose $(A, U) \neq \emptyset$. Fix an arbitrary element $* \in (A, U)$. Consider the simplicial set

$$(A, \mathcal{U}_X) : \dots \rightarrow (A, U_X^3) \rightarrow (A, U_X^2) \rightarrow (A, U).$$

The very definition of homotopy groups [GJ99, I.7.] $p_n := \pi_n((A, \mathcal{U}_X), *)$ and the universal property of the fiber product yields $p_n = 0$ for $n \geq 1$. The homotopy set p_0 is a quotient of (A, U) : morphisms $f, g : A \rightarrow U$ are identified iff there is a morphism $h : A \rightarrow U \times_X U$ such that $f = pr_1 \circ h$, $g = pr_2 \circ h$. For the simplicially constant complex $\mathcal{Q} = (p_0)_{n \geq 0}$ we obviously have $\pi_n(\mathcal{Q}) = 0$ for $n \geq 1$, $\pi_0(\mathcal{Q}) = p_0$. Thus, the canonical morphism of simplicial sets $(A, \mathcal{U}_X) \rightarrow \mathcal{Q}$ is a weak equivalence in the sense of [GJ99, I.7., page 32]. Weak equivalences are preserved by $\mathbb{Z}(-)$, thus the associated morphism of Moore complexes $[A, \mathcal{U}_X] \rightarrow \mathbb{Z}(\mathcal{Q})$ is a quasi-isomorphism [GJ99, Proposition III.2.16., Corollary III.2.7.]. So, our complex is exact at $[A, U_X^n]$, $n > 1$.

Exactness at $[A, U]$ follows from the definition of the fiber product: Let $s = \sum_{i \in I} n_i \cdot s_i$ be mapped to 0 under f . One inductively reduces to the case $\#I = 2$. Then it is obvious that there is $g \in [A, U_X^2]$, such that $g \mapsto s$. \square

H-topologies

DEFINITION 1.5. [Voe96] A morphism of schemes $f : X \rightarrow Y$ is called *topological epimorphism*, if the topological space associated to Y is a quotient topological space of X , i.e. if f is surjective and $A \subset X$ is open iff $f(A) \subset Y$ is open. f is called *universal topological epimorphism*, if this holds for any base change.

Coverings of the *h-topology* on \mathbf{Sch} are by definition finite families $\{f_i : U_i \rightarrow X\}$ of morphisms of finite type, such that $\sqcup f_i : \sqcup U_i \rightarrow X$ is a universal topological epimorphism. If the f_i are furthermore quasi-finite, this family is defined to be a covering of the *qfh-topology*.

The *cdh-topology* is the minimal Grothendieck topology containing Nisnevich coverings and abstract blow-ups (see above).

Normal qfh-coverings $\{f_i : U_i \rightarrow X\}$ are qfh-coverings having a factorization $f_i = f \circ j_i$, where $\{j_i : U_i \rightarrow U\}$ is a Zariski covering by open subsets, $f : U \rightarrow X$ a finite surjective morphism.

A family of morphisms $\{f_i : U_i \rightarrow X\}$ is called *normal h-covering*, if it has a factorization $f_i = b \circ f \circ j_i$, where $\{j_i : U_i \rightarrow U\}$ is an open covering, $f : U \rightarrow X_Z$ finite surjective and $b : X_Z \rightarrow X$ the blow-up of X of a closed center Z .

Any scheme is h-locally smooth by Proposition 1.1. We have the following inclusions of topologies on \mathbf{Sch} : $\text{Zar} \subseteq \text{Nis} \subseteq \text{ét} \subseteq \text{qfh} \subseteq \text{h}$ and $\text{Nis} \subseteq \text{cdh} \subseteq \text{h}$.

PROPOSITION 1.6. [Voe96, Theorem 3.1.9] *Every h-covering of a scheme admits a normal refinement. The same is true for qfh-coverings of smooth schemes.*

LEMMA 1.7. *Let \mathcal{A} be an abelian category closed under countable direct sums. Let $\mathcal{B} \subset \mathcal{D} := \mathbf{D}^-(\mathcal{A})$ be a full triangulated subcategory closed under existing countable direct sums (i.e. \aleph_1 -localising in the sense of Definition A.3). Then, for any complex $F \in \mathcal{D}$ with $F^n \in \mathcal{B}$ we have $F \in \mathcal{B}$.*

Proof. Compare [MVW, Lemma 9.4.]. □

We write $\mathcal{D}' := \mathbf{D}^-(\mathbf{PSh}(\mathbf{Sch}))$ and $\mathcal{D} := \mathbf{D}^-(\mathbf{Shv}_h(\mathbf{Sch}))$ in the following lemmas. The localising subcategory generated by a class of objects T , i.e. the minimal triangulated subcategory closed under existing direct sums containing T will be denoted by $\langle T \rangle$ (see Definition A.2).

LEMMA 1.8. *We have $\langle \mathbb{Z}_h(X), X \in \mathbf{Sm} \rangle = \mathcal{D}$ and $\langle \mathbb{Z}(X), X \in \mathbf{Sch} \rangle = \mathcal{D}'$.*

Proof. For any h-sheaf F , the morphism $\bigoplus_{X \in \mathbf{Sm}, a \in F(X)} \mathbb{Z}_h(X) \xrightarrow{\oplus a} F$ is surjective in $\mathbf{Shv}_h(\mathbf{Sch})$ because of weak resolution of singularities. This easily yields the assertion (Lemma 1.7). The statement for presheaves is proven similarly. □

LEMMA 1.9. *Let $T_h \subset \mathcal{D}'$ denote the class of augmented Moore complexes of h-coverings $U \rightarrow X$, where $U \in \mathbf{Sm}$, $X \in \mathbf{Sch}$ (Definition 1.2). The sheafification functor $\mathcal{D}' \rightarrow \mathcal{D}$ factorizes over $\mathcal{D}' / \langle T_h \rangle$ and induces an equivalence of categories $\mathcal{D}' / \langle T_h \rangle \cong \mathcal{D}$.*

Proof. Let \mathcal{C}_h denote the localising subcategory of \mathcal{D}' generated by presheaves F with $F_h = 0$. With Lemma 1.7, it is quite obvious to see that we have $\mathcal{D} \cong \mathcal{D}' / \mathcal{C}_h$, so it suffices to show $T_h = \mathcal{C}_h$. We have $\langle T_h \rangle \subseteq \mathcal{C}_h$ (Lemma 1.3).

For the converse, choose a complex $F \in \mathcal{C}_h$. We may assume that it is concentrated in degree 0 (Lemma 1.7). Using weak resolution of singularities and the technique of Lemma 1.8, we obtain an epimorphism of presheaves $\bigoplus_i \text{coker } \mathbb{Z}(f_i) \rightarrow F$, where $f_i : U_i \rightarrow X_i$ are h-coverings, U_i smooth. As $(\text{coker } \mathbb{Z}(f_i))_h = 0$, we even get a resolution of F by sheaves of this form. We may thus assume $F = \text{coker } f$ for an h-covering $f : U \rightarrow X$ (Lemma 1.7). The sequence

$$\dots \rightarrow \mathbb{Z}(U \times_X U) \rightarrow \mathbb{Z}(U) \rightarrow \mathbb{Z}(X) \rightarrow \text{coker } f \rightarrow 0$$

is exact (Lemma 1.4). The covering lies in $\langle T_h \rangle$ and so does $F = \text{coker } f$. □

Remark 1.10. The proof shows that the proposition still holds, if we enlarge T_h by adding complexes becoming exact when they are h-sheafified. In particular one may add arbitrary h-hypercoverings (Lemma 1.3). On the other hand, it suffices to localise with respect to h-coverings $\mathbb{Z}(\mathcal{U}_X) \rightarrow \mathbb{Z}(X)$, where $U \rightarrow X$ is of normal form (Proposition 1.6).

2. Sheaves with transfers

We are going to study the relationship between sheaves with and without transfers. It turns out that every h-sheaf has a unique transfer structure. Note that the sheafification of a presheaf with transfers is in general (i.e. for an arbitrary topology) not necessarily a presheaf with transfers.

LEMMA 2.1. *The canonical (mono-)morphisms $\mathbb{Z}(X) \rightarrow \mathbb{Z}_{\text{tr}}(X)$, $X \in \mathbf{Sm}$ induce isomorphisms*

$$\mathbb{Z}_{\text{h}}(X) \cong \mathbb{Z}_{\text{tr,h}}(X), \quad \mathbb{Q}_{\text{h}}(X) \cong \mathbb{Q}_{\text{tr,h}}(X).$$

They induce a transfer structure on any h-sheaf, which is called canonical transfer structure.

Proof. This lemma was proven independently by Ivorra [Ivo05, Corollaire 3.2.19]. The second part of the lemma was announced without proof by Voevodsky [Voe00a, §3.4].

Fix a smooth scheme V . By definition, $c(X/k, 0)(V) = c(X \times_k V/V, 0)$ is the group of universally integral cycles of $X \times_k V$, whose support is proper of relative dimension 0 over V [SV00, §2.3., 3.1.1-3.1.3, definition after Lemma 3.3.9]. This presheaf is obviously additive. Since all presheaves we are dealing with are additive, we may suppose that V is connected. To be proper of relative dimension 0 is the same as to be finite over V . As V is regular, any relative cycle is universally integral [SV00, Proposition 3.3.15]. V is normal, thus geometrically “unibranche” [Gro67, 6.15.1, 0.23.2.1], therefore any subset $Z \subseteq X$ which is equidimensional over V defines a relative cycle [SV00, Theorem 3.4.2].

$$c(X/k, 0)(V) \supset \mathbb{Z} \left[\begin{array}{l} Z \subset X \times V, \text{ finite over} \\ V, \text{ equidimensional over } V \end{array} \right]$$

Any effective cycle is equidimensional [SV00, Proposition 3.1.7]. Thus we obtain for any smooth connected V :

$$c^{\text{eff}}(X/k, 0)(V) = \mathbb{N}[Z \subseteq X \times V, Z \text{ finite over } V, Z \text{ irred.}] =: \mathbb{N}_{\text{tr}}(X)(V),$$

i.e. $c^{\text{eff}}(X/k, 0)(V) = \text{Hom}_{\mathbb{N}\mathbf{SmCor}}(V, X)$ (the free abelian monoid generated by elementary correspondences from V to X). The functoriality of the two presheaves agrees [SV00, Corollary 3.3.11]. We obtain an equality of presheaves on \mathbf{Sm} : $c^{\text{eff}}(X/k, 0)|_{\mathbf{Sm}} = \mathbb{N}_{\text{tr}}(X)$, and a fortiori $c_{\text{h}}^{\text{eff}}(X/k, 0) = \mathbb{N}_{\text{tr,h}}(X)$. Moreover, the map $\mathbb{N}(X) \rightarrow c^{\text{eff}}(X/k, 0)$ mapping a morphism of schemes to its graph induces an isomorphism $\mathbb{N}_{\text{h}}(X) \xrightarrow{\cong} c_{\text{h}}^{\text{eff}}(X/k, 0)$ [SV00, Theorem 4.2.12(2)].

Assembling these isomorphisms together we conclude that the canonical monomorphism $\mathbb{N}(X) \rightarrow \mathbb{N}_{\text{tr}}(X)$ becomes an isomorphism when we apply h-sheafification. This yields the assertions for \mathbb{Z}_{tr} and \mathbb{Q}_{tr} as well.

The canonical transfer structure is given as follows: let F be an h-sheaf (without transfers), $X \in \mathbf{Sm}$. Then we have

$$\begin{aligned} F(X) &= \text{Hom}_{\mathbf{PSh}(\mathbf{Sm})}(\mathbb{Z}(X), F) = \text{Hom}_{\mathbf{Shv}_{\text{h}}(\mathbf{Sm})}(\mathbb{Z}_{\text{h}}(X), F) \\ &\cong \text{Hom}_{\mathbf{Shv}_{\text{h}}(\mathbf{Sm})}(\mathbb{Z}_{\text{tr,h}}(X), F) = \text{Hom}_{\mathbf{PSh}(\mathbf{Sm})}(\mathbb{Z}_{\text{tr}}(X), F). \end{aligned}$$

Let $Z \in \text{Hom}_{\mathbf{SmCor}}(Y, X) = \text{Hom}_{\mathbf{PSh}_{\text{tr}}(\mathbf{Sm})}(\mathbb{Z}_{\text{tr}}(Y), \mathbb{Z}_{\text{tr}}(X))$. We define $F(Z) : F(X) \rightarrow F(Y)$ using the above isomorphism by $a \mapsto a \circ \mathbb{Z}_{\text{tr}}(Z)$. This clearly defines a transfer structure on F whose restriction to \mathbf{Sm} coincides with the original h-sheaf on \mathbf{Sm} . \square

PROPOSITION 2.2. *There is an equivalence of categories of h-sheaves with and without transfers:*

$$\mathbf{Shv}_{\text{h,tr}}(\mathbf{Sm}) \cong \mathbf{Shv}_{\text{h}}(\mathbf{Sch}).$$

Proof. Note that h-sheaf on \mathbf{SmCor} (i.e. on \mathbf{Sm}) is meant in the sense of the induced topology under the embedding $\mathbf{Sm} \subset \mathbf{Sch}$ [SGA, Exposé II, 1.1., 3.1.]. As any scheme is h-locally smooth, we have an equivalence of h-sheaves on \mathbf{Sch} and h-sheaves on \mathbf{Sm} [SGA, Exposé III, Théorème 4.1].

We make the following obvious remark: for any presheaf with transfers G the following functors on \mathbf{Sm} are naturally isomorphic:

$$G(-) \cong \mathrm{Hom}_{\mathbf{PSh}_{\mathrm{tr}}(\mathbf{Sm})}(\mathbb{Z}_{\mathrm{tr}}(-), G) \cong \mathrm{Hom}_{\mathbf{PSh}(\mathbf{Sm})}(\mathbb{Z}_{\mathrm{tr}}(-), G).$$

The first bijection is the Yoneda lemma. The second one is shown similarly using the transfer structure of G .

First, we show that for any presheaf with transfers F the canonical morphism $F \rightarrow F_{\mathrm{h}}$ is a morphism of presheaves with transfers. F_{h} has its canonical transfer structure (Lemma 2.1), in particular we have $F_{\mathrm{h}}(X) = \mathrm{Hom}_{\mathbf{PSh}(\mathbf{Sm})}(\mathbb{Z}_{\mathrm{tr}}(X), F_{\mathrm{h}})$. For a correspondence $Z \in \mathrm{Hom}_{\mathbf{SmCor}}(Y, X)$ the morphism $F(X) \rightarrow F(Y)$ (or $F_{\mathrm{h}}(X) \rightarrow F_{\mathrm{h}}(Y)$) is induced by the map $\mathbb{Z}_{\mathrm{tr}}(Y) \rightarrow \mathbb{Z}_{\mathrm{tr}}(X)$. Consider the following diagram:

$$\begin{array}{ccc} F(X) = \mathrm{Hom}_{\mathbf{PSh}(\mathbf{Sm})}(\mathbb{Z}_{\mathrm{tr}}(X), F) & \longrightarrow & \mathrm{Hom}_{\mathbf{PSh}(\mathbf{Sm})}(\mathbb{Z}_{\mathrm{tr}}(X), F_{\mathrm{h}}) = F_{\mathrm{h}}(X) \\ \downarrow & & \downarrow \\ F(Y) = \mathrm{Hom}_{\mathbf{PSh}(\mathbf{Sm})}(\mathbb{Z}_{\mathrm{tr}}(Y), F) & \longrightarrow & \mathrm{Hom}_{\mathbf{PSh}(\mathbf{Sm})}(\mathbb{Z}_{\mathrm{tr}}(Y), F_{\mathrm{h}}) = F_{\mathrm{h}}(Y) \end{array}$$

The horizontal maps are induced by morphisms of presheaves (*without* transfers) $F \rightarrow F_{\mathrm{h}}$. The commutativity of the diagram yields the first claim.

Secondly, one analogously proves that morphisms of h-sheaves are morphisms of h-sheaves with transfers.

Thirdly, we show that any h-sheaf admits only one transfer structure. Considering any h-sheaf F , the canonical morphism $F \rightarrow F_{\mathrm{h}}$ is an isomorphism of presheaves without transfer. According to the first claim, it is also a morphism of sheaves with transfers, where F_{h} carries the canonical transfer structure. This implies that F carries the canonical transfer structure, as well. \square

Cohomology comparison results

Recall that a presheaf F on \mathbf{Sch} (or \mathbf{Sm}) is called *homotopy invariant*, if the canonical projection $p : X \times \mathbb{A}^1 \rightarrow X$ induces an isomorphism $p^* : F(X) \rightarrow F(X \times \mathbb{A}^1)$ for any X in \mathbf{Sch} (or \mathbf{Sm} , respectively). One easily sees that for any complex F^* of homotopy invariant presheaves the cohomology presheaves $H^n(F^*)$ are homotopy invariant, as well.

First of all, we gather several results of Friedlander, Suslin and Voevodsky in the following theorem. These facts constitute the nontrivial input to our comparison result:

THEOREM 2.3. *Let k be a field admitting strong resolutions of singularities. Let F be an homotopy invariant presheaf of \mathbb{Q} -vector spaces with transfers. Let t be either the Zariski, Nisnevich, étale or cdh-topology. Then the cohomology presheaves on \mathbf{Sm}*

$$X \mapsto H_t^n(X, F_t), n \geq 0$$

are homotopy invariant presheaves with transfers and are (up to canonical isomorphisms) independent of t .

Proof. Presheaves with transfers are pretheories [Voe00b, Proposition 3.1.11]. Hence [Voe00a, Theorem 4.27, Proposition 5.28] (Zariski-topology), [Voe00a, Theorem 5.7.] (Nisnevich), [Voe00a, Propositions 5.24, 5.27] (étale topology, this requires rational coefficients) and finally the proof of [FV00, Theorem 5.5] (cdh-topology, this requires strong resolution of singularities) apply. \square

We want to generalize this theorem to the qfh- and h-topology. We first observe that the h-topology is the “union” of the cdh- and the qfh-topology, in the sense made precise by Lemma 2.4. Looking at the definitions of these topologies and Proposition 1.6, this statement is not surprising.

We then use the usual Galois descent method in order to overcome the gap between Nisnevich and qfh-topology (Lemma 2.6). The cohomology comparison result between étale and qfh-topology (for a qfh-sheaf, Theorem 2.7) and the adaptation of the cohomology comparison result of Nisnevich vs. cdh-topology to the case qfh- vs. h-topology then give the result.

LEMMA 2.4. *Let F be a presheaf on \mathbf{Sch} . We assume that F is a qfh-sheaf and a cdh-sheaf. Then F is an h-sheaf.*

Proof. Considering the exact sequence of qfh-sheaves

$$0 \longrightarrow \ker \longrightarrow F \longrightarrow F_h \longrightarrow \text{coker} \longrightarrow 0,$$

we are reduced to show $F = 0$ for any F with $F_h = 0$.

Let $X \in \mathbf{Sch}$, without loss of generality X is reduced (as F is a qfh-sheaf), $a \in F(X)$. There is an h-covering $\{g_i : U_i \rightarrow X\}$, such that $g_i^*(a) = 0$. We may assume that the covering is of normal form (Proposition 1.6):

$$g_i : U_i \xrightarrow{f_i} U \xrightarrow{f} X_{X_1} \xrightarrow{p} X.$$

Clearly $\{U_i \xrightarrow{f \circ f_i} X_{X_1}\}$ is a qfh-covering. Hence $p^*(a) = 0$, for F is a qfh-sheaf. Since F is a cdh-sheaf and $X_{X_1} \sqcup X_1 \rightarrow X$ is a cdh-covering, it suffices to show $F(X_1) = 0$.

We proceed by noetherian induction with respect to the strictly descending chain of closed subsets $X =: X_0 \supsetneq X_1 \supsetneq \dots$. The induction stops when the chain becomes stationary or if the normal refinement of the covering of X_i does not contain any blow-up. The former is impossible, thus at some point of the noetherian induction the h-covering of X_i must even be a qfh-covering. \square

THEOREM 2.5. *(Comparison of Nisnevich and (qf)h-sheafification)*

We assume that k admits strong resolution of singularities. Let F be a homotopy invariant presheaf of \mathbb{Q} -vector spaces with transfers (on \mathbf{Sm}). Then the Nisnevich, qfh- and h-sheafifications F_{Nis} , $F_{\text{qfh}}|_{\mathbf{Sm}}$ and $F_h|_{\mathbf{Sm}}$ are isomorphic.

Consider the presheaves $\mathbb{Q}_{\text{tr}'}(X)$ [SV96, §5 and §6], where $X \in \mathbf{Sch}$:

$$\mathbb{Q}_{\text{tr}'}(X)(Y) := \mathbb{Q} \left[\begin{array}{l} W \subset X \times Y, \text{ closed, integral,} \\ \text{finite and surjective over } Y \end{array} \right],$$

where Y is an arbitrary normal connected scheme. Analogously we have a presheaf $\mathbb{Z}[1/p]_{\text{tr}'}(X)$, where p is the exponential characteristic of k . Functoriality is given as follows: Let $f : Z \rightarrow Y$ be a morphism of normal connected schemes, $W \subset X \times Y$ an integral cycle, finite and surjective over Y . Set $f^*([W]) := \sum n_i [(V_i)_{\text{red}}]$, where V_i are the irreducible components of $W \times_Y Z$ and $(V_i)_{\text{red}}$ are the associated reduced (thus integral) schemes. The multiplicities n_i are defined as described in [SV96, definition after Lemma 5.6]. They are always positive, see *loc. cit.*

LEMMA 2.6. *Let $f : Y \rightarrow X$ be the normalization of a smooth (thus excellent) connected affine scheme X in a normal finite field extension L of $k(X)$. Then there exists a correspondence*

$$f_* \in \mathbb{Z} \left[\frac{1}{p} \right]_{\text{tr}'}(Y)(X)$$

(finite integral correspondence from X to Y with coefficients in $\mathbb{Z}[1/p]$), such that $f \circ f_ = d \cdot \text{Id}_X$, $d \in \mathbb{N}^{>0}$.*

Proof. (compare [Voe96, Theorem 3.3.8] for \mathbb{Z}_{qfh}).

The automorphism group $G := \text{Aut}(Y/X) = \text{Gal}(k(Y)/k(X))$ is a finite group. Considering $g \in G$ as the graph of the corresponding automorphism, the sum $a = \sum_{g \in G} g$ is a G -invariant

element of $\mathbb{Z}_{\text{tr}'}(Y)(Y)$ (a is an integral cycle as Y is integral). According to [SV96, Corollary 6.2], a has a preimage $f_* \in \mathbb{Z}[1/p]_{\text{tr}'}(Y)(X)$:

$$\begin{array}{ccccc} Y & & & & \\ \downarrow f & \searrow a & & & \\ X & \xrightarrow{f_*} & Y & \xrightarrow{f} & X \end{array}$$

Let d denote the cardinality of G . Then $f \circ f_* \circ f = f \circ a = d \cdot f$.

We show that for any surjective morphism $f : Y \rightarrow X$ of schemes, where Y is assumed to be integral, and X to be smooth and every effective correspondence $c \in \mathbb{N}_{\text{tr}'}(X)(X)$ with $c \circ f = d \cdot f$ the equality $c = d \cdot \text{Id}_X$ holds. This provides the assertion of the lemma, for f is surjective in our situation and – by the proof of *loc. cit.* – f_* , being the preimage of an effective correspondence, is effective as well.

Let $c = \sum_j n_j \cdot c_j$, $n_j \in \mathbb{N}$, $c_j \subset X \times X$ integral and finite over X . By definition

$$c \circ f = \sum_{i,j} n_j \cdot m_{ij} \cdot W_{ij} \stackrel{!}{=} d \cdot \Gamma_f,$$

where $W_{ij} \subset T_j$ are the associated reduced schemes of the irreducible components of $T_j := c_j \times_X Y$, the morphism $c_j \rightarrow X$ is the projection to the first component. The multiplicities of the components m_{ij} are positive (see above). As all coefficients are positive, it follows $W_{ij} = \Gamma_f$ for all i, j and $\sum_{i,j} n_j m_{ij} = d$.

Using the surjectivity of f it is easy to see (using Hilbert's Nullstellensatz) that the \bar{k} -valued points of Δ_X and c_j agree. Thus $\Delta_X = c_j$ on the level of topological spaces (Hilbert's Nullstellensatz). As X , and thus Δ_X is reduced, they agree as schemes, too. Hence we have $\#\{j\} = 1$ and $\#\{i\} = 1$ as $T = \Gamma_f \cong Y$ is irreducible. Moreover $m = 1$, for Y is integral by assumption. \square

Proof of Theorem 2.5. In view of Theorem 2.3 we may assume that F is an homotopy invariant cdh-sheaf with transfers, which implies that F_{qfh} is an h-sheaf (Lemma 2.4), i.e. $F_{\text{qfh}} = F_{\text{h}}$. Consider the exact sequence of Nisnevich sheaves with transfers:

$$0 \longrightarrow F' \longrightarrow F \longrightarrow F_{\text{qfh}} \longrightarrow F'' \longrightarrow 0.$$

(Recall that $\mathbf{Shv}_{\text{Nis, tr}}(\mathbf{Sm})$ is an abelian category [Voe00b, Theorem 3.1.4] and that $F \rightarrow F_{\text{qfh}} = F_{\text{h}}$ is a morphism of sheaves with transfers, see Proposition 2.2).

As in the proof of Lemma 2.4 it suffices to show the following: Let F be a (not necessarily homotopy invariant) cdh-sheaf on \mathbf{Sch} , whose restriction to \mathbf{Sm} is a sheaf with transfers, such that $F_{\text{qfh}} = 0$. Then we have $F = 0$.

Let $X \in \mathbf{Sch}$, without loss of generality affine and smooth (as F is a cdh-sheaf), and $a \in F(X)$ an arbitrary section. We show $a = 0$. There is a qfh-covering of normal form $\{f_i : U_i \rightarrow U \rightarrow X\}$, such that $f_i^*(a) = 0$ (Proposition 1.6). As F is a Zariski sheaf, we may suppose that our covering consists of a single surjective morphism. We replace the excellent scheme U by its normalization in a normal finite extension of $k(X)$. Choose a cdh-covering $\tilde{U} \rightarrow U$ with $\tilde{U} \in \mathbf{Sm}$ (strong resolution of singularities).

As F is a sheaf with transfers, $F(V) = \text{Hom}_{\mathbf{PSh}_{\text{tr}}(\mathbf{Sm})}(\mathbb{Q}_{\text{tr}}(V), F) = \text{Hom}_{\mathbf{PSh}(\mathbf{Sm})}(\mathbb{Q}_{\text{tr}}(V), F) = \text{Hom}_{\mathbf{PSh}(\mathbf{Sm})}(\mathbb{Q}_{\text{tr}'}(V), F)$ for any smooth V (see the proof of Proposition 2.2). As F is a cdh-sheaf, we may cdh-sheafify the first presheaf in all of these Hom-sets without changing the Hom-sets. Hence, the morphism $F(X) \rightarrow F(\tilde{U})$ is induced by

$$\mathbb{Q}_{\text{tr}', \text{cdh}}(\tilde{U}) \xrightarrow{\alpha} \mathbb{Q}_{\text{tr}', \text{cdh}}(U) \xrightarrow{\beta} \mathbb{Q}_{\text{tr}', \text{cdh}}(X).$$

We know $\mathbb{Q}_{\text{tr}'}(W)|_{\mathbf{Sm}} = \mathbb{Q}_{\text{qfh}}(W)|_{\mathbf{Sm}}$ [SV96, Theorem 6.3] for any scheme W , a fortiori $\mathbb{Q}_{\text{tr}'}(W)_{\text{cdh}} = \mathbb{Q}_{\text{qfh,cdh}}(W) = \mathbb{Q}_{\text{h}}(W)$. As $\tilde{U} \rightarrow U$ is an h-covering, α is an epimorphism [Voe96, Proposition 3.2.5.2]. According to Lemma 2.6, β has a section, so it is an epimorphism. Hence $F(X) \rightarrow F(\tilde{U})$ is a monomorphism, i.e. $a = 0$. \square

THEOREM 2.7. *(Comparison of étale and qfh-cohomology) We assume that k admits strong resolution of singularities. Let F be a homotopy invariant étale sheaf with transfers of \mathbb{Q} -vector spaces and let X be a normal scheme. Then $H_{\text{ét}}^n(X, F) \cong H_{\text{qfh}}^n(X, F_{\text{qfh}})$, $n \in \mathbb{N}$.*

Proof. The case $n = 0$ is Theorem 2.5, so we may assume that F is a qfh-sheaf. Then [Voe96, Theorem 3.4.1] is what we assert. \square

THEOREM 2.8. *(Comparison of h- and qfh-cohomology) Let k be a field admitting strong resolution of singularities and let F be an homotopy invariant presheaf with transfers of \mathbb{Q} -vector spaces. Then for all $X \in \mathbf{Sm}$, $n \geq 0$ the following holds:*

$$H_{\text{h}}^n(X, F_{\text{h}}) = H_{\text{qfh}}^n(X, F_{\text{qfh}}).$$

LEMMA 2.9. *Let $X \in \mathbf{Sm}$, $Z \subset X$ a smooth closed subscheme. Let $p_Z : X_Z \rightarrow X$ denote the blow-up of Z in X . Further, let F be a homotopy invariant presheaf with transfers of \mathbb{Q} -vector spaces. Then it holds for $i \geq 0$*

$$\text{Ext}_{\mathbf{Shv}_{\text{qfh}}(\mathbf{Sm})}^i(\text{coker } \mathbb{Z}_{\text{qfh}}(p_Z), F_{\text{qfh}}) = 0.$$

Proof. An analogous statement for the Nisnevich topology is proven in [Voe00a, Proposition 5.21] exploiting the fact that Nisnevich cohomology groups of F_{Nis} are homotopy invariant. The proof of *loc. cit.* remains valid for any finer topology t , which fulfills $H_t^i(X, F_t) \cong H_t^i(X \times \mathbb{A}^1, F_t)$. In particular it holds (for sheaves of \mathbb{Q} -vector spaces) for the qfh-topology (Theorem 2.3, Theorem 2.7). \square

LEMMA 2.10. *Let k be a field admitting strong resolution of singularities, F a qfh-sheaf, such that $F_{\text{h}} = 0$, G a homotopy invariant presheaf with transfers of \mathbb{Q} -vector spaces. Then we have*

$$\text{Ext}_{\mathbf{Shv}_{\text{qfh}}(\mathbf{Sm})}^n(F, G_{\text{qfh}}) = 0, n \geq 0.$$

Proof. The proof of [FV00, Lemma 5.4] goes through (using Lemma 2.9 instead of [FV00, Lemma 5.3]), if we replace Nisnevich and cdh-topology by qfh- and h-topology respectively. \square

Proof of Theorem 2.8. The proof of [FV00, Theorem 5.5] goes through verbatim using Lemma 2.10 instead of [FV00, Lemma 5.4]. \square

The following theorem gathers the above results. It points out the quite special nature of homotopy invariant Zariski sheaves with transfers of \mathbb{Q} -vector spaces: they are already sheaves with respect to the (much finer) h-topology.

THEOREM 2.11. *Let $t \in \{\text{Zar}, \text{Nis}, \text{ét}, \text{cdh}, \text{qfh}, \text{h}\}$. Let F^* be a bounded above complex of presheaves of \mathbb{Q} -vector spaces with transfers, whose cohomology- t -sheaves are homotopy invariant. We assume that k admits strong resolution of singularities. Then the hypercohomology presheaves $X \mapsto \mathbb{H}_t^n(X, F_t^*)$, $X \in \mathbf{Sm}$ are homotopy invariant presheaves with transfers and are independent of the choice of the topology t .*

The assumption on F^ is satisfied, if the cohomology presheaves of F^* are homotopy invariant.*

Proof. If F^* is concentrated in degree 0, we are done because of Theorem 2.3, Theorem 2.7, Theorem 2.5, Theorem 2.8, as hypercohomology is just cohomology in this case. This also shows the last claim.

In general, let denote $(H^q(F^*))_t$ the t -sheafification of the q -th cohomology presheaf of the complex F^* , which is an homotopy invariant presheaf with transfers. The presheaves $X \mapsto \mathbb{H}_{\text{Nis}}^n(X, F_{\text{Nis}}^*)$

have transfers [MVW, Exercise 13.5.]. So, the other hypercohomology groups, which are canonically isomorphic to the Nisnevich hypercohomology, have transfers as well. The hypercohomology spectral sequence [Wei94, Proposition 5.7.9]

$$E_2^{p,q} = H_t^p(X, (H^q(F^*))_t) = \text{Ext}^p(\mathbb{Q}_t(X), (H^q(F^*))_t) \implies \mathbb{H}_t^{p+q}(X, F_t^*)$$

converges because of finite cohomological dimension for sheaves of \mathbb{Q} -vector spaces (see [Voe96, Theorem 3.4.6] for the h-topology). Comparing the limit terms for X and $X \times \mathbb{A}^1$ yields the assertion. \square

Remark 2.12. We restrict to \mathbb{Q} -linear categories, as the comparison of Nisnevich and qfh-sheaves is valid only for sheaves of \mathbb{Q} -vector spaces. An analogous statement for sheaves with torsion does not hold: Consider the Nisnevich sheaf with transfers \mathcal{O}^* . If R is a ring, then $\mathcal{O}^*(R[t]) = R^* \times \{1 + \sum_{i=1}^n f_i t^i, (f_i)^N = 0, N \gg 0\}$. In particular, \mathcal{O}^* is homotopy invariant on \mathbf{Sm} , but not on \mathbf{Sch} . See [Voe00a, Remark 5.8] for a counterexample for the Nisnevich-flat topology.

The restriction to characteristic zero is only needed for the cohomology comparison result of Nis- and cdh-topology (and the analogue qfh- vs. h-topology). It seems reasonable to aim at a generalization to positive characteristic at least for \mathbb{Q} -coefficients using Noether normalization, but we did not succeed to prove it.

3. Motivic complexes and geometrical motives

3.1 Motivic complexes

As we have outlined in the introduction, we will embed the several categories of geometrical motives – which we are mostly interested in – into the more flexible categories of motivic complexes. Voevodsky defines a category of motivic complexes using Nisnevich-sheaves with transfers. We will also use a different version using h-sheaves without transfers. In the case of \mathbb{Q} -vector space sheaves we will show that both categories are equivalent (Theorem 3.7). This theorem relies on the cohomology comparison results of Chapter 2.

Recall from Definition A.2 that $\langle - \rangle$ denotes the minimal localising subcategory containing a given class of objects.

DEFINITION 3.1. [Voe00b, §3.1.] The category of *effective motivic complexes with transfers* is defined to be the full subcategory $\mathbf{DM}_{\text{Nis,tr}}^{\text{eff},-}$ of $\mathbf{D}^-(\mathbf{Shv}_{\text{Nis,tr}}(\mathbf{Sm}))$ consisting of complexes whose Nisnevich cohomology sheaves are homotopy invariant. (This category is denoted by $\mathbf{DM}_{\text{Nis}}^{\text{eff},-}(k)$ in *loc. cit.*).

Similarly, $\mathbf{DM}_{\text{h}}^{\text{eff},-} \subset \mathbf{D}^-(\mathbf{Shv}_{\text{h}}(\mathbf{Sch}))$ denotes the full subcategory, whose objects have homotopy invariant cohomology h-sheaves. This category is called category of *effective motivic complexes without transfers*.

We write $\mathbf{DM}_{\text{h}}^{\text{eff},-}(\mathbb{Q})$ and $\mathbf{DM}_{\text{Nis,tr}}^{\text{eff},-}(\mathbb{Q})$ for the corresponding \mathbb{Q} -linear categories.

DEFINITION 3.2. Let Δ^* denote the standard cosimplicial object in \mathbf{Sch} , i.e. $\Delta^n = \mathbb{A}^{n+1} / (\sum x_i = 1)$. For any presheaf F , let $C_*(F)$ denote the Suslin-complex $F(\Delta^* \times -)$, i.e. $C_n(F)(X) = F(\Delta^n \times X)$. Using the Dold-Kan-construction we obtain a complex of presheaves $C_*(F)$ (whose boundary homomorphisms are alternating sums of δ_j) and thus an exact functor $C_* : \mathbf{PSh}(\mathbf{Sch}) \rightarrow \mathbf{Kom}(\mathbf{PSh}(\mathbf{Sch}))$. If F is a sheaf with respect to an arbitrary (pre)topology, then $C_*(F)$ is a complex of sheaves as well as well.

For the rest of this subsection, we will use the following notation: $\mathcal{D}_{\text{h}} := \mathbf{D}^-(\mathbf{Shv}_{\text{h}}(\mathbb{Q}\mathbf{Sch}))$, $\mathcal{A}_{\text{h}} := \langle \{\mathbb{Q}_{\text{h}}(X \times \mathbb{A}^1) \rightarrow \mathbb{Q}_{\text{h}}(X), X \in \mathbf{Sm}\} \rangle \subset \mathcal{D}_{\text{h}}$.

Remark 3.3. Morphisms whose cone is in \mathcal{A}_h are called *weak \mathbb{A}^1 -equivalences*. Objects $L \in \mathcal{D}_h$ are called *\mathbb{A}^1 -local*, if for all weak \mathbb{A}^1 -equivalences $K \rightarrow K'$ the induced map $\mathrm{Hom}(K', L) \rightarrow \mathrm{Hom}(K, L)$ is a bijection.

Theorem 2.11 implies that $\mathbf{DM}_h^{\mathrm{eff}, -}(\mathbb{Q})$ is a triangulated subcategory of \mathcal{D}_h : For a distinguished triangle $A \rightarrow B \rightarrow C \rightarrow A[1]$ in \mathcal{D}_h with $A, B \in \mathbf{DM}_h^{\mathrm{eff}, -}$ we have that $\mathrm{Hom}_{\mathcal{D}_h}(\mathbb{Q}_h(-), A[n]) = \mathbb{H}_h^n(-, A)$ is homotopy invariant, analogously for B . The long exact $\mathrm{Hom}_{\mathcal{D}_h}$ -sequence coming from this triangle shows that C is \mathbb{A}^1 -local as well, compare [MVW, Lemma 9.17.]. Hence, C has homotopy invariant cohomology sheaves. The latter can be seen as in [MVW, Proposition 14.5] (using Theorem 2.11). As in *loc. cit.* one sees that bounded above complexes (of sheaves of \mathbb{Q} -vector spaces) with homotopy invariant cohomology sheaves are \mathbb{A}^1 -local.

PROPOSITION 3.4. *With these notations there is an equivalence of categories $\mathcal{D}_h/\mathcal{A}_h \cong \mathbf{DM}_h^{\mathrm{eff}, -}(\mathbb{Q})$ induced by a triangulated functor $\mathbf{RC} : \mathcal{D}_h \rightarrow \mathbf{DM}_h^{\mathrm{eff}, -}(\mathbb{Q})$, whose restriction on $\mathbf{Shv}_h(\mathbf{Sch})$ is given by C_* .*

Proof. (compare [Voe00b, Proposition 3.2.3]). For brevity, we omit the specification of \mathbb{Q} -vector space sheaves in the notations. We show that the restriction of the projection map $\Phi : \mathbf{DM}_h^{\mathrm{eff}, -} \subset \mathcal{D}_h \xrightarrow{\mathrm{Proj.}} \mathcal{D}_h/\mathcal{A}_h$ is an equivalence of categories and that the restriction to h-sheaves of the inverse equivalence Φ^{-1} is just C_* :

$$\begin{array}{ccc} \mathbf{Shv}_h(\mathbf{Sch}) & \longrightarrow & \mathcal{D}_h/\mathcal{A}_h \\ & \searrow C_* & \downarrow \Phi^{-1} \\ & & \mathbf{DM}_h^{\mathrm{eff}, -} \end{array}$$

Then, we set $\mathbf{RC} := \Phi^{-1} \circ \mathrm{Proj.}$ According to Remark 3.3 and [Nee01, Theorem 2.1.8], Φ is triangulated and so is \mathbf{RC} .

Firstly, we show that Φ is fully faithful. Secondly, we show the essential surjectivity. Both parts will use the cohomology comparison results for homotopy invariant sheaves with transfers.

Φ is fully faithful: Let $T \in \mathbf{DM}_h^{\mathrm{eff}, -}$ be arbitrary. According to Proposition 2.2, $T|_{\mathbf{Sm}}$ is a complex of sheaves with transfers, which implies (Theorem 2.11) for all $X \in \mathbf{Sm}, n \in \mathbb{Z}$

$$\mathrm{Hom}_{\mathcal{D}_h}(\mathbb{Q}_h(X), T[n]) \cong \mathrm{Hom}_{\mathcal{D}_h}(\mathbb{Q}_h(X \times \mathbb{A}^1), T[n]).$$

Hence (compare [MVW, Lemma 9.17]) T is an \mathbb{A}^1 -local object in \mathcal{D}_h . Then we have for all $S \in \mathcal{D}_h$: $\mathrm{Hom}_{\mathcal{D}_h/\mathcal{A}_h}(S, T) \cong \mathrm{Hom}_{\mathcal{D}_h}(S, T)$ [Nee01, Lemma 9.1.5].

Φ is essentially surjective: we know $K \cong \mathrm{Tot}(C_*K)$ in $\mathcal{D}_h/\mathcal{A}_h$ (see [MVW, Lemma 9.14], the proof without transfers is strictly parallel, as the homotopy providing the quasi-isomorphism is induced by the multiplication map $\mu : \mathbb{A}^2 \rightarrow \mathbb{A}^1$). According to Proposition 2.2, K and $\mathrm{Tot}(C_*(K))$ are complexes of h-sheaves with transfers (of \mathbb{Q} -vector spaces). The cohomology presheaves of $\mathrm{Tot}(C_*(K))$ are homotopy invariant [MVW, Lemma 2.17], thus the associated h-sheaves of the cohomology presheaves restricted to \mathbf{Sm} are homotopy invariant, as well (Theorem 2.11). Using the equivalence of h-sheaves on \mathbf{Sm} and on \mathbf{Sch} (see the proof of Proposition 2.2), this easily implies that the same holds without restricting to \mathbf{Sm} . Hence $\mathrm{Tot}(C_*(K)) \in \mathbf{DM}_h^{\mathrm{eff}, -}$ and we are done. \square

Remark 3.5. If one attempts to define explicitly $\Phi^{-1} : K \mapsto \mathrm{Tot}(C_*K)$, one has to deal with the question, whether C_* is an exact functor. It suffices to show the right-exactness of C_* . The presheaf cokernel F of an epimorphism of h-sheaves has transfers (Proposition 2.2) and satisfies $F_h = 0$. Using the cohomology comparison result for the h-topology (Theorem 2.11) instead of the Nisnevich analogue [MVW, Theorem 13.11.] yields that the h-sheafification of $C_*(F)$ is exact. The h-analogue of [MVW, Corollary 13.13.] shows that C_* is well-defined on \mathcal{D}_h .

LEMMA 3.6. *The categories $\mathbf{D}^-(\mathbf{Shv}_h(\mathbb{Q}\mathbf{Sch}))$ and $\mathbf{DM}_h^{\mathrm{eff}, -}(\mathbb{Q})$ are tensor triangulated categories.*

Proof. Starting with $X \otimes Y := X \times_k Y$, $X, Y \in \mathbf{Sch}$, one obtains a tensor structure on $\mathbf{D}^-(\mathbf{Shv}_h(\mathbf{Sch}))$ in the same manner as in [MVW, Corollary 8.17].

The tensor structure on $\mathcal{D}_h/\mathcal{A}_h$ is determined by the requirement that the canonical functor $\mathcal{D}_h \rightarrow \mathcal{D}_h \rightarrow \mathcal{A}_h$ is a tensor functor, compare [MVW, Corollary 9.4.]. Proposition 3.4, i.e. $\mathcal{D}_h/\mathcal{A}_h \cong \mathbf{DM}_h^{\text{eff},-}(\mathbb{Q})$ yields the second assertion. \square

THEOREM 3.7. *Let k be a field admitting strong resolution of singularities. Using the notations of Definition 3.1 there are natural equivalences of tensor triangulated categories*

$$\mathbf{DM}_{\text{Nis,tr}}^{\text{eff},-}(\mathbb{Q}) \cong \mathbf{DM}_h(\mathbb{Q}) \cong \mathbf{DM}_h^{\text{eff},-}(\mathbb{Q}),$$

where $\mathbf{DM}_h(\mathbb{Q})$ denotes the homological category of h -sheaves (with rational coefficients) [Voe96, §4.1.].

Proof. The first comparison is stated (without proof) by [Voe00b, Theorem 4.1.12]. The theorem is essentially a corollary of the cohomology comparison result of Nisnevich and h -topology (Theorem 2.11) and the fact that h -sheafs have a unique transfer structure (Proposition 2.2). We permit ourselves to omit “ \mathbb{Q} ” in the notations.

Let us write $\mathcal{A}_{\text{Nis,tr}} \subset \mathcal{D}_{\text{Nis,tr}} := \mathbf{D}^-(\mathbf{Shv}_{\text{Nis,tr}}(\mathbb{Q}\mathbf{Sm}))$ for the localising subcategory generated by $\mathbb{Q}_{\text{tr}}(X \times \mathbb{A}^1) \rightarrow \mathbb{Q}_{\text{tr}}(X)$, $X \in \mathbf{Sm}$. We have an equivalence of categories $\mathbf{DM}_{\text{Nis,tr}}^{\text{eff},-} \cong \mathcal{D}_{\text{Nis,tr}}/\mathcal{A}_{\text{Nis,tr}}$ [Voe00b, Proposition 3.2.3]. We show that the composition $(-)_h : \mathbf{Shv}_{\text{Nis,tr}}(\mathbf{Sm}) \rightarrow \mathbf{Shv}_{\text{Nis}}(\mathbf{Sm}) \rightarrow \mathbf{Shv}_h(\mathbf{Sch})$ of the forgetful and sheafification functor, which clearly extends to the corresponding derived categories, maps $\mathcal{A}_{\text{Nis,tr}}$ to $0 \in \mathbf{DM}_h$. It suffices to show that the complexes generating $\mathcal{A}_{\text{Nis,tr}}$ are mapped to 0, which is equivalent to $K_X := \text{Ker}(\mathbb{Q}_{\text{tr}}(X \times \mathbb{A}^1) \rightarrow \mathbb{Q}_{\text{tr}}(X)) \mapsto 0$, i.e. that $(K_X)_h$ is contractible in the sense of [Voe96, Definition 2.2.1]. (According to Lemma 2.1, $\mathbb{Q}_{\text{tr}}(X)$, $X \in \mathbf{Sm}$ maps to $\mathbb{Q}_h(X)$). The latter holds obviously, as $K_X = \mathbb{Q}_{\text{tr}}(X) \otimes K_{\text{Spec } k}$ and $(K_{\text{Spec } k})_h$ is contractible [Voe96, Lemma 2.2.3]. We obtain a surjective functor $\Phi : \mathcal{D}_{\text{Nis,tr}}/\mathcal{A}_{\text{Nis,tr}} \rightarrow \mathbf{DM}_h$ (Proposition 2.2). (*nota bene:* We have not yet used rational coefficients).

$$\begin{array}{ccc} \mathcal{D}_{\text{Nis,tr}} & \xrightarrow{(-)_h} & \mathcal{D}_h \\ \downarrow & & \downarrow \\ \mathcal{D}_{\text{Nis,tr}}/\mathcal{A}_{\text{Nis,tr}} & \xrightarrow{\Phi} & \mathbf{DM}_h \end{array}$$

For the full faithfulness of Φ it suffices to show

$$\text{Hom}_{\mathcal{D}_{\text{Nis,tr}}}(\mathbb{Q}_{\text{tr}}(X), F[n]) = \text{Hom}_{\mathbf{DM}_h}(\mathbb{Q}_h(X), F_h[n])$$

for every $F \in \mathbf{DM}_{\text{Nis,tr}}^{\text{eff},-}$.

For brevity, we write $\text{Spec } k$ for $\mathbb{Q}_h(\text{Spec } k)$, X for $\mathbb{Q}_h(X)$ etc. The triangle $I^1 \rightarrow \mathbb{A}^1 \rightarrow \text{Spec } k \rightarrow I^1[1]$ is distinguished in \mathcal{D}_h . As the category is tensor triangulated (Lemma 3.6), we obtain a long exact sequence

$$\dots \leftarrow \text{Hom}(X \otimes I^1, F_h[n]) \leftarrow \text{Hom}(X \otimes \mathbb{A}^1, F_h[n]) \leftarrow \text{Hom}(X, F_h[n]) \leftarrow \dots$$

The two terms to the right are the n -th hypercohomologies of F in X and $X \times \mathbb{A}^1$, respectively. They are homotopy invariant (Theorem 2.11). Hence we have $\text{Hom}(X \otimes I^1, F_h[n]) = 0$, i.e. F_h is strictly homotopy invariant in the sense of [Voe96, definition 2.2.8], which implies $\text{Hom}_{\mathbf{DM}_h}(X, F_h[n]) = \text{Hom}_{\mathcal{D}_h}(X, F_h[n]) = \mathbb{H}_h^n(X, F_h)$ [Voe96, Proposition 2.2.9].

The equivalence $\mathbf{DM}_h^{\text{eff},-} \cong \mathbf{DM}_h$ is trivial (and does not need rational coefficients). \square

3.2 Geometrical motives

We now define the categories of (quasi-)geometrical motives with and without transfers. The prefix “quasi” indicates that we also consider countable direct sums (i.e. disjoint unions) of schemes. Moreover, the “quasi”-categories consist of bounded above complexes, which should be considered still being of geometric origin, but in general not being compact.

Recall from Definition A.3 that for a class T of objects of a triangulated category the notations $\langle T \rangle^{\aleph_0}$ (and $\langle T \rangle^{\aleph_1}$, respectively) mean the minimal thick subcategory containing T (closed under countable existing direct sums, respectively).

DEFINITION 3.8. The categories $\overline{\mathbb{Z}\mathbf{Sch}}$ and $\overline{\mathbf{SmCor}}$ are defined as the closure under countable direct sums of $\mathbb{Z} : \mathbb{Z}\mathbf{Sch} \rightarrow \mathbf{PSh}(\mathbf{Sch})$ (the Yoneda embedding), and similarly for $\overline{\mathbf{SmCor}}$. For brevity, we may write $\oplus X_i := \oplus \mathbb{Z}(X_i)$ for a family of schemes X_i .

DEFINITION 3.9. [Voe00b, §2.1.] Consider the class T consisting of the following complexes

- $X \times \mathbb{A}^1 \rightarrow X$, $X \in \mathbf{Sm}$.
- $U \cap V \xrightarrow{\iota_U \oplus \iota_V} U \oplus V \xrightarrow{\iota_U \oplus (-\iota_V)} X$ for any open covering $X = U \cup V$ of a smooth scheme X .

The category of *effective quasigeometric motives with transfers* $\mathbf{DM}_{\text{gm, tr}}^{\text{eff}}$ is defined as follows: $\mathbf{DM}_{\text{gm, tr}}^{\text{eff}} := \mathbf{K}^- (\overline{\mathbf{SmCor}}) / \langle T \rangle^{\aleph_1}$, i.e. the localisation of $\mathbf{K}^- (\overline{\mathbf{SmCor}})$ with respect to the minimal triangulated subcategory closed under countable sums containing T . The subcategory $\mathbf{DM}_{\text{gm, tr}}^{\text{eff}}$ consisting of $\mathbf{K}^b(\mathbf{SmCor})$ is called category of *effective geometric motives with transfers*. (This coincides with Voevodsky’s definition $\mathbf{K}^b(\mathbf{SmCor}) / \langle T \rangle^{\aleph_0}$, see Theorem 3.15).

DEFINITION 3.10. We define the category $\mathbf{DM}_{\text{gm}}^{\text{eff}}$ of *effective quasigeometric motives without transfers* over a field k : Let T denote the union of the following two classes of complexes in $\mathbf{K}^- (\overline{\mathbb{Z}\mathbf{Sch}})$:

- $T_{\mathbb{A}^1} = \{X \times \mathbb{A}^1 \xrightarrow{pr_1} X, X \in \mathbf{Sch}\}$.
- $T_{\text{h}} =$ augmented Moore-complexes of h-coverings $U \rightarrow X$, $U \in \mathbf{Sm}$, $X \in \mathbf{Sch}$ (Definition 1.2).

We define $\mathbf{DM}_{\text{gm}}^{\text{eff}}$ to be the localisation of $\mathbf{K}^- (\overline{\mathbb{Z}\mathbf{Sch}})$ with respect to the \aleph_1 -localising subcategory generated by T :

$$\mathbf{DM}_{\text{gm}}^{\text{eff}} := \mathbf{K}^- (\overline{\mathbb{Z}\mathbf{Sch}}) / \langle T \rangle^{\aleph_1}.$$

The subcategory $\mathbf{DM}_{\text{gm}}^{\text{eff}}$ of bounded complexes of schemes is called category of *effective geometric motives without transfers*. The canonical functor $M_{\text{gm}} : \mathbf{Sch} \rightarrow \mathbf{DM}_{\text{gm}}^{\text{eff}}$ maps a scheme to its *geometric motive*.

If we consider rational coefficients instead of integral ones, we write $\mathbf{DM}_{\text{gm}}^{\text{eff}}(\mathbb{Q})$ etc.

Remark 3.11. Note that the quasigeometric categories of motives are pseudo-abelian, as they are closed under direct sums of the form $\oplus_{i \in \mathbb{N}} A$, where $A \in \mathbf{DM}_{\text{gm, tr}}^{\text{eff}}$ [Lev98, Lemma II.2.2.4.8.1].

LEMMA 3.12. *The category $\mathbf{DM}_{\text{gm}}^{\text{eff}}$ is a triangulated tensor category, whose tensor structure is induced by $M_{\text{gm}}(X) \otimes M_{\text{gm}}(Y) = M_{\text{gm}}(X \times Y)$. The subcategory $\mathbf{DM}_{\text{gm}}^{\text{eff}}$ is closed under this tensor product.*

Proof. The tensor structure on $\mathbf{K}^- (\overline{\mathbb{Z}\mathbf{Sch}})$ is determined via the one on \mathbf{Sch} ($X \otimes Y := X \times Y$) and $(\oplus X_i) \otimes (\oplus Y_j) := \oplus_{i,j} X_i \otimes Y_j$ and $K^* \otimes L^* := \text{Tot}((K^n \otimes L^m)^{n,m})$. We show that for all $A \in T$ (Definition 3.10) and $B \in \mathbf{K}^- (\overline{\mathbb{Q}\mathbf{Sch}})$ the tensor product $A \otimes B$ lies in $\langle T \rangle^{\aleph_1}$. Then [Lev98, II.2.3.4] yields the claim. The subcategory $\mathcal{B} \subset \mathbf{K}^- (\overline{\mathbb{Q}\mathbf{Sch}})$ containing the complexes B with $A \otimes B \in \langle T \rangle^{\aleph_1}$ is \aleph_1 -localising. Using Lemma 1.7 we reduce to the case $B \in \mathbf{Sch}$, which is obvious. The second claim is clear from the definitions. \square

Remark 3.13. A similar statement holds for the variants with transfers $\mathbf{DM}_{\text{gm, tr}}^{\text{eff}}$ (proof as above) and $\mathbf{DM}_{\text{gm, tr}}^{\text{eff}}$ [Voe00b, Proposition 2.1.3].

3.3 Embedding theorems

THEOREM 3.14. *Let k admit the strong resolution of singularities. Then we have a commutative diagram of tensor-triangulated functors:*

$$\begin{array}{ccc} \mathbf{K}^- (\overline{\mathbf{QSch}}) & \xrightarrow{\mathbb{Q}_h} & \mathbf{D}^- (\mathbf{Shv}_h (\mathbf{QSch})) \\ \downarrow & & \downarrow \mathbf{RC} \\ \mathbf{DM}_{\text{gm}}^{\text{eff}} & \xrightarrow[\subset]{i} & \mathbf{DM}_h^{\text{eff}, -} (\mathbb{Q}) \end{array}$$

where i is a full embedding of quasigeometric effective motives (Definition 3.10) into the category of effective motivic complexes (Definition 3.1). The image of i is precisely the subcategory of \aleph_1 -compact objects of $\mathbf{DM}_h^{\text{eff}, -} (\mathbb{Q})$. The idempotent completion of the image of $\mathbf{K}^b (\mathbf{QSch})$ is the subcategory of compact objects in $\mathbf{DM}_h^{\text{eff}, -} (\mathbb{Q})$.

Proof. (compare [Voe00b, Theorem 3.2.6]). In the proof, we write $\mathcal{D}_h := \mathbf{D}^- (\mathbf{Shv}_h (\mathbf{QSch}))$ and $\mathcal{A}_h := \langle \mathbb{A}^1\text{-complexes} \rangle$.

Existence of i : By its very definition it is clear that the functor \mathbf{RC} commutes with direct sums. We show that $\mathbf{RC}(\mathbb{Q}_h(T_{\mathbb{A}^1})) = 0$ and $\mathbf{RC}(\mathbb{Q}_h(T_h)) = 0$, which yields the existence of i . The first claim directly follows from the construction of \mathbf{RC} (see Proposition 3.4, $\mathcal{A}_h = \mathbb{Q}_h(T_{\mathbb{A}^1})$). $\mathbf{RC} : \mathcal{D} \rightarrow \mathcal{D}/\mathcal{A}_h \xrightarrow{\Phi^{-1}} \mathbf{DM}_h^{\text{eff}, -}$. For the second claim, we note $\mathbf{RC}(\mathbb{Q}_h(Y)) = C_*(\mathbb{Q}_h(Y)) =: C_*(Y)$. The inverse equivalence Φ^{-1} of Proposition 3.4 maps a complex K of sheaves to the total complex $\text{Tot}(C_*(K))$. We have to show that the total complex of the double complex

$$C_*(\mathcal{U}_X \rightarrow X \rightarrow 0) : \dots \rightarrow C_*(U \times_X U) \rightarrow C_*(U) \rightarrow C_*(X) \rightarrow 0$$

is exact for any h -covering $U \rightarrow X$, $X \in \mathbf{Sch}$. But $C_*(\mathcal{U}_X \rightarrow X \rightarrow 0)$ is exact in view of the exactness of C_* (see the proof of Proposition 3.4) and Lemma 1.3. As Tot preserves quasi-isomorphisms, we are done.

Full faithfulness of i : Let us denote $\mathcal{D}' := \mathbf{D}^- (\mathbf{PSh}(\mathbf{QSch}))$. Consider the following diagram

$$\begin{array}{ccc} \mathbf{K}^- (\overline{\mathbf{QSch}}) & \xrightarrow{\mathbb{Q}(-)} & \mathcal{D}' \\ & \searrow \mathbb{Q}_h(-) & \downarrow \\ & & \mathcal{D} \stackrel{1.9}{\cong} \mathcal{D}' / \langle \mathbb{Q}(T_h) \rangle \\ & & \downarrow \mathbf{RC} \\ \mathbf{K}^- (\overline{\mathbf{QSch}}) / \langle T_{\mathbb{A}^1}, T_h \rangle^{\aleph_1} & \longrightarrow & \mathcal{D} / \langle \mathbb{Q}(T_{\mathbb{A}^1}) \rangle \cong \mathbf{DM}_h^{\text{eff}, -} \end{array}$$

The functor $\mathbb{Q} : \mathbf{K}^- (\overline{\mathbf{QSch}}) \rightarrow \mathcal{D}'$ is a full embedding [Wei94, Corollary 10.4.7] as the presheaves $\oplus_i \mathbb{Q}(X_i)$ are direct sums of projective objects, thus projective. Proposition A.5 precisely grants the extension of the embedding $\mathbf{K}^- (\overline{\mathbf{QSch}}) \subset \mathcal{D}'$ to the categories in question and the asserted compactness statements.

i is tensor-triangulated: The functors \mathbf{RC} and \mathbb{Q}_h are tensor functors by definition. It is clear that \mathbb{Q}_h is tensor-triangulated. In order to see that \mathbf{RC} is tensor-triangulated, it suffices to remark that the projection $\mathcal{D} \rightarrow \mathcal{D} / \langle \mathbb{Q}_h(T_{\mathbb{A}^1}) \rangle$ is tensor-triangulated. Hence, $\mathbf{RC} \circ \mathbb{Q}_h$ is tensor-triangulated, which yields the claim. \square

THEOREM 3.15. *There is a full embedding*

$$i_{\text{tr}} : \mathbf{DM}_{\text{gm},\text{tr}}^{\text{eff}} \subset \mathbf{DM}_{\text{Nis},\text{tr}}^{\text{eff},-}.$$

Similarly to the theorem above, its essential image is exactly the subcategory of \aleph_1 -compact objects and the idempotent completion of $i_{\text{tr}}(\mathbf{K}^b(\mathbf{SmCor}))$ are the compact objects of $\mathbf{DM}_{\text{Nis},\text{tr}}^{\text{eff},-}$.

Proof. The proof of [Voe00b, Theorem 3.2.6] generalizes to the quasi-geometric situation (using the \aleph_1 -compactness arguments as in Theorem 3.14, Remark A.6). \square

4. Reinterpretation of geometrical motives

We are now ready to gather the results of the preceding sections in order to achieve the main result of the present paper:

THEOREM 4.1. *Let k be a field admitting strong resolution of singularities. Then there is an equivalence of tensor triangulated categories between the categories of effective geometric motives (with rational coefficients) with and without transfers (Definitions 3.9, 3.10):*

$$\mathbf{DM}_{\text{gm}}^{\text{eff}}(\mathbb{Q}) \cong \mathbf{DM}_{\text{gm},\text{tr}}^{\text{eff}}(\mathbb{Q}).$$

The same is true for the quasigeometric categories:

$$\mathbf{DM}_{\text{gm}}^{\text{eff}}(\mathbb{Q}) \cong \mathbf{DM}_{\text{gm},\text{tr}}^{\text{eff}}(\mathbb{Q}).$$

The theorem simplifies the handling of geometric motives insofar as one does not have to work with correspondences. Moreover, singular schemes are contained in the category in a natural manner. Last but not least, the inversion of the h-topology permits greater flexibility in the manipulation of objects. The price for this flexibility is the loss of torsion information.

Proof. We embed the categories of effective (quasi)-geometric motives into the categories of motivic complexes, see Theorems 3.14, 3.15. Either in the case with and without transfers, these are exactly the (\aleph_1) -compact objects (Definition A.3, Proposition A.5). The equivalence $\mathbf{DM}_{\text{h}}^{\text{eff},-}(\mathbb{Q}) \cong \mathbf{DM}_{\text{Nis},\text{tr}}^{\text{eff},-}(\mathbb{Q})$ (Theorem 3.7) yields the assertion. \square

Remark 4.2. An attempt to prove an analogous statement for a category of geometric motives defined by inverting only smooth h-hypercoverings fails, because the Moore complex of a hypercovering is in general not exact at the level of presheaves.

At first glance, the equivalence of the geometric categories, i.e. bounded complexes might be surprising, as a non-smooth scheme is *ad hoc* only replaced by an unbounded smooth h-hypercovering. We describe a bounded complexes of smooth schemes isomorphic to a scheme X : there is a proper surjective morphism $p : \tilde{X} \rightarrow X$, which is an isomorphism outside some closed subset $Z \subsetneq X$ (strong resolution of singularities). We obtain a distinguished triangle in $\mathbf{DM}_{\text{gm},\text{tr}}^{\text{eff}}$ of the form

$$M_{\text{gm}}(p^{-1}(Z)) \rightarrow M_{\text{gm}}(Z) \oplus M_{\text{gm}}(\tilde{X}) \rightarrow M_{\text{gm}}(X) \rightarrow M_{\text{gm}}(p^{-1}(Z))[1]$$

[Voe96, Proposition 4.1.5]. Inductively applying this resolution technique to Z and $p^{-1}(Z)$ one constructs a so-called hypercube of schemes, one of whose vertices is X , the other vertices are smooth. According to the above exact triangle, the motive of X is isomorphic to the total complex of the rest of the hypercube. The induction stops after at most $\dim X$ steps. See [GNAPGP88, especially Theorem 2.15] for the construction of cubical hyperresolutions.

In the sequel, we will draw some corollaries of the theorem. We will always assume that k admits strong resolution of singularities and we will work with rational coefficients.

COROLLARY 4.3. *If A, B are closed subschemes of a scheme X , such that $A \cup B = X$, then there is a distinguished triangle in $\mathbf{DM}_{\text{gm}}^{\text{eff}}$ of the following form:*

$$M_{\text{gm}}(A \cap B) \rightarrow M_{\text{gm}}(A) \oplus M_{\text{gm}}(B) \rightarrow M_{\text{gm}}(X) \rightarrow M_{\text{gm}}(A \cap B)[1].$$

In particular, as a motive, a scheme is isomorphic to the associated reduced scheme: $M_{\text{gm}}(X_{\text{red}}) \cong M_{\text{gm}}(X)$.

Proof. This follows from the fact that $A \sqcup B \rightarrow X$ is a qfh-covering and Lemma 1.4. \square

DEFINITION 4.4. Inverting the Tate-Object $\mathbb{Q}(1)$ in the category of effective geometric motives (with transfers), Voevodsky defines the category of geometrical motives, which we denote by $\mathbf{DM}_{\text{gm},\text{tr}}$ [Voe00b, §2.1, page 192]. Strictly parallel, we define the category of *geometrical motives (without transfers)* \mathbf{DM}_{gm} and their quasi-geometric analogs $\mathbf{DM}_{\overline{\text{gm}}}$ and $\mathbf{DM}_{\overline{\text{gm}},\text{tr}}$.

THEOREM 4.5. *The above equivalence of effective geometrical motives induces an equivalence of geometrical motives with and without transfers (with rational coefficients):*

$$\mathbf{DM}_{\text{gm}}(\mathbb{Q}) \cong \mathbf{DM}_{\text{gm},\text{tr}}(\mathbb{Q}).$$

Moreover, the canonical functor $\mathbf{DM}_{\text{gm}}^{\text{eff}}(\mathbb{Q}) \rightarrow \mathbf{DM}_{\text{gm}}(\mathbb{Q})$ is a full embedding. These statements are also true for the quasigeometrical categories.

Proof. As $\mathbf{DM}_{\text{gm}}^{\text{eff}}(\mathbb{Q}) \cong \mathbf{DM}_{\text{gm},\text{tr}}^{\text{eff}}(\mathbb{Q})$ is an equivalence of tensor categories, it maps $\mathbb{Q}(n) \in \mathbf{DM}_{\text{gm}}^{\text{eff}}(\mathbb{Q})$ to $\mathbb{Q}(n) \in \mathbf{DM}_{\text{gm},\text{tr}}^{\text{eff}}(\mathbb{Q})$ (note that the equivalence restricted to complexes of smooth schemes is the identity). So, the first statement in either the geometric or the quasi-geometric case is clear from Theorem 4.1. The second statement is then a reformulation of [Voe00b, Theorem 4.3.1] in the geometric case. The quasi-geometric case follows from the proof of *loc. cit.* and the fact that \otimes commutes with \oplus . \square

The reinterpretation of geometrical motives may also be used to get a more conceptual proof of results of Huber concerning realizations of motives [Hub00, Hub04]:

COROLLARY 4.6. *Let \mathcal{A} be any \mathbb{Q} -linear abelian category. Let $R : \mathbb{Q}\text{Sch} \rightarrow \mathbf{D}(\mathcal{A})$ be a functor mapping Čech nerves of h -coverings and \mathbb{A}^1 -complexes (see Definition 3.10) to complexes quasi-isomorphic to zero. Then it extends to a functor $\mathbf{DM}_{\text{gm}}^{\text{eff}} \rightarrow \mathbf{D}(\mathcal{A})$. If furthermore \mathcal{A} is a tensor category and tensoring in \mathcal{A} with $R(\mathbb{Q}(1))$ is an equivalence of categories, R passes to $\mathbf{DM}_{\overline{\text{gm}}} \rightarrow \mathbf{D}(\mathcal{A})$. In particular, singular realization of varieties factors over geometrical motives.*

Appendix A. Compact objects in categories of bounded above complexes

In order to motivate the somewhat lengthy definition of \aleph_1 -compact objects, we first outline the application of these notions. It turns out that the abstract characterization of complexes of schemes in terms of compactness notions is quite useful – note the simplicity of the proof of Theorem 4.1. Consider the following special case of a theorem of Neeman [Nee01, Theorem 4.4.9], which is (tacitly) used in the proof of the embedding theorem of Voevodsky [Voe00b, Theorem 3.2.6]:

PROPOSITION A.1. *The full embedding $\mathbf{K}^b(\mathbb{Z}\text{Sch}) \subset \mathbf{D}(\mathbf{PSh}(\text{Sch}))$ identifies the former category with the subcategory of compact objects of the latter category. Localizing with respect to some compact objects, for example Mayer-Vietoris complexes $U \cap V \rightarrow U \oplus V \rightarrow U \cup V$ the above embedding extends to an embedding*

$$\mathbf{K}^b(\mathbb{Z}\text{Sch})/M.-V. \subset \mathbf{D}(\mathbf{PSh}(\text{Sch}))/M.-V.$$

Note that the right hand side is just $\mathbf{D}(\mathbf{Shv}_{\text{Zar}}(\mathbf{Sch}))$. Instead of the Zariski topology, we want to invert the h-topology. In this setting, we cannot replace the unbounded Moore complexes of h-coverings by some bounded complexes. Hence we are looking for a similar statement for $\mathbf{K}^- (\mathbb{Z}\mathbf{Sch})$ instead of $\mathbf{K}^b (\mathbb{Z}\mathbf{Sch})$. In order to apply the aforementioned theorem of Neeman, we enlarge $\mathbf{K}^- (\mathbb{Z}\mathbf{Sch})$ insofar as we add countable existing direct sums; we arrive at $\mathbf{K}^- (\overline{\mathbb{Z}\mathbf{Sch}})$, which is precisely the subcategory of so-called \aleph_1 -compact objects (see below). Then a similar statement as above holds (see Proposition A.5 for the precise statement).

Neeman theorem handles the case that the larger category is closed under arbitrary direct sums. So, we will have to make some minor modifications to Neeman's definitions in order to adapt them to bounded above categories like $\mathbf{D}^-(\mathbf{PSh}(\mathbf{Sch}))$. The idea is to add boundedness conditions in all definitions and to check that the proof of [Nee01, Theorem 4.4.9] remains valid.

In the following definitions, let β be an infinite cardinal, let \mathcal{A} be an abelian category having all direct sums, e.g. $\mathcal{A} = \mathbf{PSh}(\mathbf{Sch})$, and $\mathcal{D} := \mathbf{D}^-(\mathcal{A})$.

DEFINITION A.2. A triangulated subcategory $\mathcal{S} \subset \mathcal{D}$ is called $(\beta\text{-})$ localising, if it is closed under direct sums (of cardinality $< \beta$) that exist in \mathcal{D} . The smallest thick $(\beta\text{-})$ localising subcategory containing a class of objects S will be denoted $\langle S \rangle$ ($\langle S \rangle^\beta$, respectively). Hence $\langle S \rangle^{\aleph_0}$ is the minimal thick subcategory containing S and $\langle S \rangle^{\aleph_1}$ is the minimal subcategory closed under countable existing direct sums. We only need these two cases.

To any complex $C^* \in \mathcal{D}$, we associate an integer $|C| := \max\{n, C^n \neq 0\}$. (Recall that our complexes have differentials of degree $+1$). Let $\{X_\lambda, \lambda \in \Lambda\}$ be a set of objects in \mathcal{D} . It is called bounded if $\sup\{|X_\lambda|, \lambda \in \Lambda\}$ is finite, which implies that the direct sum $\bigoplus X_\lambda$ exists (in \mathcal{D}). We make the following definitions (note that working in the category $\mathbf{D}(\mathcal{A})$ and dropping the boundedness conditions, we exactly arrive at Neeman's definitions [Nee01, §§3,4]).

DEFINITION A.3. – A class S of objects of \mathcal{D} is called β -perfect if it contains the zero object and the following conditions hold:

1. Let $\{X_\lambda, \lambda \in \Lambda\}$ be a bounded collection of cardinality $< \beta$ of objects in \mathcal{D} . Any morphism $k \rightarrow \bigoplus X_\lambda$ with $k \in S$ factors as $k \rightarrow \bigoplus k_\lambda \xrightarrow{\oplus f_\lambda} \bigoplus X_\lambda$, where $k_\lambda \in S$. Furthermore, if Λ is infinite, we also require $\sup |k_\lambda| \leq \sup |X_\lambda|$ (which ensures that $\bigoplus k_\lambda$ exists).
 2. If any such morphism $k \rightarrow \bigoplus k_\lambda \xrightarrow{\oplus f_\lambda} \bigoplus X_\lambda$ vanishes, then there are objects $l_\lambda \in S$ such that f_λ is factorized by $k_\lambda \rightarrow l_\lambda \rightarrow X_\lambda$ and $k \rightarrow \bigoplus k_\lambda \rightarrow \bigoplus l_\lambda$ vanishes. Again, if Λ is an infinite set, we require $\sup |l_\lambda| \leq \sup |X_\lambda|$.
- There is a maximal β -perfect class of objects, contained in \mathcal{S} [Nee01, Corollary 3.3.10]. It will be denoted \mathcal{S}_β .
 - An object $k \in \mathcal{D}$ is called β -small, if any morphism $k \rightarrow \bigoplus_{\lambda \in \Lambda} X_\lambda$, where the family X_λ is supposed to be bounded, factors as $k \rightarrow \bigoplus_{\lambda \in \Lambda'} X_\lambda \rightarrow \bigoplus_{\lambda \in \Lambda} X_\lambda$, where $\Lambda' \subset \Lambda$ is a subset of cardinality $< \beta$. The full subcategory of \mathcal{D} consisting of β -small objects is denoted $\mathcal{D}^{(\beta)}$.
 - The subcategory of β -compact objects in \mathcal{D} is defined by $\mathcal{D}^\beta := \{\mathcal{D}^{(\beta)}\}_\beta$. If $\beta = \aleph_0$, we will usually speak of compact objects.

LEMMA A.4. The compact objects of $\mathcal{D}' := \mathbf{D}^-(\mathbf{PSh}(\mathbf{Sch}))$ are given by the subcategory (under the Yoneda embedding) $\mathbf{K}^b(\mathbb{Z}\mathbf{Sch})$ of bounded complexes of schemes. The \aleph_1 -compact objects of \mathcal{D}' are precisely given by bounded above complexes of countable direct products of schemes, i.e. $\mathbf{K}^- (\overline{\mathbb{Z}\mathbf{Sch}})$. The same is true for $\mathcal{D} := \mathbf{D}^-(\mathbf{Shv}_h(\mathbf{Sch}))$.

Proof. A scheme, considered as an object of \mathcal{D}' under the Yoneda embedding, is clearly \aleph_0 -small (i.e. $\text{Hom}_{\mathcal{D}'}(\mathbb{Z}(X), -)$ commutes with direct sums). The same is true in the category \mathcal{D} . In fact, this holds for any topology t whose coverings are open morphisms: by convention our schemes are

quasi-compact, so every covering has a finite subcovering, which implies that the direct sum of a set of t -sheaves F_λ is given by $X \mapsto \bigoplus F_\lambda(X)$. Via induction on the length of a complex one sees that a bounded complex of schemes is \aleph_0 -small. All \aleph_0 -small objects are (\aleph_0) -compact [Nee01, Example 3.3.16]. As the subcategory of β -compact objects is β -localising for a regular cardinal β (such as \aleph_0 or \aleph_1) [Nee01, Lemma 4.2.5] countable direct sums of schemes are \aleph_1 -compact. Bounded above complexes of such objects are \aleph_1 -compact as well (Lemma 1.7). We saw in Lemma 1.8 that $\langle \mathbf{K}^b(\mathbb{Z}\mathbf{Sch}) \rangle = \mathcal{D}'$, so [Nee01, Lemma 4.4.5] shows $\mathbf{K}^b(\mathbb{Z}\mathbf{Sch}) = \mathcal{D}'^{\aleph_0}$ and $\mathbf{K}^- (\overline{\mathbb{Z}\mathbf{Sch}}) = \mathcal{D}'^{\aleph_1}$ and similarly for \mathcal{D} . \square

PROPOSITION A.5. We write $T_h := \{\mathbb{Q}(\mathcal{U}_X) \rightarrow \mathbb{Q}(X), U \rightarrow X \text{ an } h\text{-covering}\}$ (see Definition 1.2) and $T_{\mathbb{A}^1} = \{\mathbb{Q}(X \times \mathbb{A}^1) \rightarrow \mathbb{Q}(X), X \in \mathbf{Sm}\}$.

The natural functor $\mathbf{K}^- (\overline{\mathbb{Q}\mathbf{Sch}}) / \langle T_h \rangle^{\aleph_1} \rightarrow \mathcal{D} := \mathbf{D}^- (\mathbf{Shv}_h(\mathbb{Q}\mathbf{Sch}))$ is a full embedding whose image consists of the \aleph_1 -compact objects of \mathcal{D} .

The natural functor

$$\mathbf{K}^- (\overline{\mathbb{Q}\mathbf{Sch}}) / \langle T_h, T_{\mathbb{A}^1} \rangle^{\aleph_1} \rightarrow \mathcal{D} / \langle T_{\mathbb{A}^1} \rangle$$

is a full embedding. Its image consists of the \aleph_1 -compact objects of \mathcal{D} . Moreover, the idempotent completion of the image of bounded complexes of schemes are precisely the compact objects of \mathcal{D} .

Proof. For the convenience of the reader, we indicate the necessary changes to the proof of [Nee01, Theorem 4.4.9]: The second part of the proof [Nee01, Lemma 3.3.3] does not work, but this lemma is only applied to triangulated subcategories, so the second part is not needed in view of [Nee01, Lemma 3.3.5]. [Nee01, Lemma 3.3.7] only holds for $\alpha \leq \beta$ or $\beta > \aleph_0$ in the notations of *loc. cit.* This lemma is only used in [Nee01, Theorem 3.3.9], which is only needed with $\alpha = \beta$. [Nee01, Theorem 4.3.3] has to be stated as follows: Let β be a regular cardinal, S a class of objects in \mathcal{D}^β . Let $f : x \rightarrow z$ be a morphism with $x \in \mathcal{D}^\beta$ and $z \in \langle S \rangle$. Then there is an object $y \in \langle S \rangle^\beta$, such that $|y| \leq |z|$ and f factors as $x \rightarrow y \rightarrow z$. Its proof is the same as in *loc. cit.* This sharper statement is necessary to prove [Nee01, Lemma 4.4.8] in our situation. The proof of the fact that localisation preserves direct sums [Nee01, Lemma 3.2.10] does à priori not work in general, but when we localise with respect to h -coverings, the claim holds, as schemes are quasi-compact (cf. the proof of Lemma A.4). When we localise with respect to \mathbb{A}^1 -complexes, the assertion is true because of Proposition 3.4.

The first claim of our proposition is now precisely the statement of [Nee01, Theorem 4.4.9]: $\mathbf{K}^- (\overline{\mathbb{Z}\mathbf{Sch}})$ are the \aleph_1 -compact objects in $\mathbf{D}^- (\mathbf{PSh}(\mathbf{Sch}))$, which is generated by these complexes and \mathcal{D} is the localisation of the \mathcal{D}' with respect to $\langle T_h \rangle$ (Lemma 1.9). The second claim follows from the first one (using [Nee01, Theorem 4.4.9] again). \square

Remark A.6. The above proposition also holds for Nisnevich sheaves with transfers. Indeed, by [Voe00b, Proposition 3.1.9] $\mathrm{Hom}_{\mathbf{D}^- (\mathbf{Shv}_{\mathrm{Nis}, \mathrm{tr}}(\mathbf{Sm}))} (X, K[i]) = \mathbb{H}_{\mathrm{Nis}}^i(X, K)$, Nisnevich hypercohomology commutes with direct sums and Proposition 3.4 is replaced by its predecessor [Voe00b, Proposition 3.2.3].

REFERENCES

- AKMW02 Dan Abramovich, Kalle Karu, Kenji Matsuki, and Jaroslaw Włodarczyk, *Torification and factorization of birational maps*, J. Amer. Math. Soc. **15** (2002), no. 3, 531–572 (electronic).
- Bon02 Laurent Bonavero, *Factorisation faible des applications birationnelles (d’après Abramovich, Karu, Matsuki, Włodarczyk et Morelli)*, Astérisque (2002), no. 282, Exp. No. 880, vii, 1–37, Séminaire Bourbaki, Vol. 2000/2001.
- dJ96 A. J. de Jong, *Smoothness, semi-stability and alterations*, Inst. Hautes Études Sci. Publ. Math. (1996), no. 83, 51–93.

JAKOB SCHOLBACH

- Ful98 William Fulton, *Intersection theory*, second ed., Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics, vol. 2, Springer-Verlag, Berlin, 1998.
- FV00 Eric M. Friedlander and Vladimir Voevodsky, *Bivariant cycle cohomology*, Cycles, transfers, and motivic homology theories, Ann. of Math. Stud., vol. 143, Princeton Univ. Press, Princeton, NJ, 2000, pp. 138–187.
- GJ99 Paul G. Goerss and John F. Jardine, *Simplicial homotopy theory*, Progress in Mathematics, vol. 174, Birkhäuser Verlag, Basel, 1999.
- GNAPGP88 F. Guillén, V. Navarro Aznar, P. Pascual Gainza, and F. Puerta, *Hyperrésolutions cubiques et descente cohomologique*, Lecture Notes in Mathematics, vol. 1335, Springer-Verlag, Berlin, 1988.
- Gro67 A. Grothendieck, *Éléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas IV*, Inst. Hautes Études Sci. Publ. Math. (1964 - 1967), no. 20, 24, 28, 32, 361.
- GS96 Henri Gillet and Christophe Soulé, *Descent, motives and K-theory*, J. Reine Angew. Math. **478** (1996), 127–176.
- Hir64 Heisuke Hironaka, *Resolution of singularities of an algebraic variety over a field of characteristic zero. I, II*, Ann. of Math. (2) **79** (1964), 109–203; *ibid.* (2) **79** (1964), 205–326.
- Hir75 ———, *Flattening theorem in complex-analytic geometry*, Amer. J. Math. **97** (1975), 503–547.
- Hub00 Annette Huber, *Realization of Voevodsky’s motives*, J. Algebraic Geom. **9** (2000), no. 4, 755–799.
- Hub04 A. Huber, *Corrigendum to: “Realization of Voevodsky’s motives”*, J. Algebraic Geom. **13** (2004), no. 1, 195–207.
- Ivo05 Florian Ivorra, *Réalisation l-adique des motifs mixtes*, 2005, Thesis, Université Paris 6, <http://www.math.uiuc.edu/K-theory/0762>.
- Lev98 Marc Levine, *Mixed motives*, Mathematical Surveys and Monographs, vol. 57, American Mathematical Society, Providence, RI, 1998.
- MVW Carlo Mazza, Vladimir Voevodsky, and Charles Weibel, *Lectures in Motivic Cohomology*, Preprint, <http://www.math.rutgers.edu/~weibel/motiviclectures.html>.
- Nee01 Amnon Neeman, *Triangulated categories*, Annals of Mathematics Studies, vol. 148, Princeton University Press, Princeton, NJ, 2001.
- RG71 Michel Raynaud and Laurent Gruson, *Critères de platitude et de projectivité. Techniques de “platification” d’un module*, Invent. Math. **13** (1971), 1–89.
- Sch05 Jakob Scholbach, *Reinterpretation geometrischer Motive in der h-Topologie*, 2005, <http://www.fmi.uni-leipzig.de/~huber/preprints.html>.
- SGA *Théorie des topos et cohomologie étale des schémas. Tome 1: Théorie des topos*, Séminaire de Géométrie Algébrique du Bois-Marie 1963–1964 (SGA 4).
- SGA72 *Théorie des topos et cohomologie étale des schémas. Tome 2*, Springer-Verlag, Berlin, 1972, Séminaire de Géométrie Algébrique du Bois-Marie 1963–1964 (SGA 4), Dirigé par M. Artin, A. Grothendieck et J. L. Verdier. Avec la collaboration de N. Bourbaki, P. Deligne et B. Saint-Donat, Lecture Notes in Mathematics, Vol. 270.
- SV96 Andrei Suslin and Vladimir Voevodsky, *Singular homology of abstract algebraic varieties*, Invent. Math. **123** (1996), no. 1, 61–94.
- SV00 ———, *Relative cycles and Chow sheaves*, Cycles, transfers, and motivic homology theories, Ann. of Math. Stud., vol. 143, Princeton Univ. Press, Princeton, NJ, 2000, pp. 10–86.
- Voe96 V. Voevodsky, *Homology of schemes I*, Selecta Mathematica **2** (1996), no. 1, 111 – 153.
- Voe00a Vladimir Voevodsky, *Cohomological theory of presheaves with transfers*, Cycles, transfers, and motivic homology theories, Ann. of Math. Stud., vol. 143, Princeton Univ. Press, Princeton, NJ, 2000, pp. 87–137.

GEOMETRIC MOTIVES AND THE H-TOPOLOGY

- Voe00b ———, *Triangulated categories of motives over a field*, Cycles, transfers, and motivic homology theories, Ann. of Math. Stud., vol. 143, Princeton Univ. Press, Princeton, NJ, 2000, pp. 188–238.
- VSF00 Vladimir Voevodsky, Andrei Suslin, and Eric M. Friedlander, *Cycles, transfers, and motivic homology theories*, Annals of Mathematics Studies, vol. 143, Princeton University Press, Princeton, NJ, 2000.
- Wei94 Charles A. Weibel, *An introduction to homological algebra*, Cambridge Studies in Advanced Mathematics, vol. 38, Cambridge University Press, Cambridge, 1994.

Jakob Scholbach scholbach@math.uni-leipzig.de

Universität Leipzig, Mathematisches Institut, Augustusplatz 10/11, 04105 Leipzig, Germany