

Multiplication of convex sets in $C(K)$ spaces

by

JOSÉ PEDRO MORENO (Madrid) and ROLF SCHNEIDER (Freiburg)

Abstract. Let $C(K)$ denote the Banach algebra of continuous real functions, with the supremum norm, on a compact Hausdorff space K . For two subsets of $C(K)$, one can define their product by pointwise multiplication, just as the Minkowski sum of the sets is defined by pointwise addition. Our main interest is in correlations between properties of the product of closed order intervals in $C(K)$ and properties of the underlying space K . When K is finite, the product of two intervals in $C(K)$ is always an interval. Surprisingly, the converse of this result is true for a wide class of compacta. We show that a first-countable space K is finite whenever it has the property that the product of two nonnegative intervals is closed, or the property that the product of an interval with itself is convex. That some assumption on K is needed can be seen from the fact that if K is the Stone–Čech compactification of \mathbb{N} , then the product of two intervals in $C(K)$ with continuous boundary functions is always an interval. For any K , it is proved that the product of two positive intervals is always an interval, and that the product of two nonnegative intervals is always convex. Finally, square roots of intervals are investigated, with results of similar type.

1. Introduction. Relations between properties of the Banach space $C(K)$ of continuous real functions on a compact Hausdorff space and properties of the underlying space K have been an object of study in analysis and topology for decades. Vector addition of closed convex sets in a real vector space, also called Minkowski addition, is a much investigated operation with many applications. These two facts were a motivation to initiate in [11] a study of Minkowski addition of closed convex sets in $C(K)$. We restricted ourselves there to particularly simple classes of convex sets, mainly intervals and intersections of balls. An *interval* $[f, g]$ in $C(K)$ is the set of all $h \in C(K)$ with $f \leq h \leq g$, where f and g are bounded real functions on K (not necessarily elements of $C(K)$). Intervals in this sense are closed and convex, and the set of intervals is closed under Minkowski addition (a nontrivial fact). One of the aims of [11] was to investigate to what extent familiar properties

2010 *Mathematics Subject Classification*: Primary 52A05; Secondary 46J10, 46E15.

Key words and phrases: $C(K)$ space, order interval, product of sets.

Received 14 January 2016; revised 7 March 2016.

Published online *.

of Minkowski addition in Euclidean spaces carry over to intervals and other convex sets in $C(K)$. As it turned out, some of these properties characterize the underlying space K as being extremally disconnected.

Now, $C(K)$ is not only a Banach space, but a Banach algebra, and this fact suggests studying not only pointwise addition, but also pointwise multiplication of convex sets. That multiplication of sets in $C(K)$, even of open balls, bears some surprises and difficulties was already evident from the papers [2–5, 9, 10]. In [9], for example, it is proved that multiplication in $C(K)$ is an open mapping if and only if K has topological (covering) dimension zero.

In the following, we study mainly multiplication of intervals in $C(K)$. Let us first mention that in \mathbb{R} (that is, when K is a singleton) the algebraic product of two compact intervals I, J , defined by $I \cdot J := \{ab : a \in I, b \in J\}$, is again an interval. In \mathbb{R}^n (that is, when K is finite) the same is true if we replace intervals by Cartesian products of intervals. Therefore, if the product of two subsets $\emptyset \neq A, B \subset C(K)$ is defined by $A \cdot B = \{h\eta : h \in A, \eta \in B\}$, it is natural to ask whether for two intervals $[f, g], [\varphi, \psi]$ their product $[f, g] \cdot [\varphi, \psi]$ is an interval. This is true, for example, if $f > 0$ and $\varphi > 0$ (Thm. 3.2). Perhaps surprisingly, it already becomes false if, say, $f > 0$ and $\varphi \geq 0$. More precisely, if K is first-countable and if the latter condition, together with continuity of f, g, φ, ψ , is sufficient for $[f, g] \cdot [\varphi, \psi]$ to be closed, then K must be finite (Thm. 4.1).

The condition that K be first-countable is needed to ensure, when K is infinite, the existence of a nontrivial convergent sequence, a technical tool used in our constructions. That some assumption of this kind is necessary is made evident by the fact that in $C(\beta\mathbb{N})$ (where $\beta\mathbb{N}$ denotes the Stone–Čech compactification of the discrete space \mathbb{N} of natural numbers) the product of two intervals $[f, g], [\varphi, \psi]$ with continuous functions f, g, φ, ψ is always an interval (Prop. 4.2).

While the product of intervals in $C(K)$ for general K remains unexplored at present, we exhibit more instances where first-countable spaces K with desirable properties of the product of intervals in $C(K)$ must be finite (thus, the exceptional role of extremally disconnected spaces in [11] is here played by finite spaces). While the product of two nonnegative intervals in $C(K)$ is always convex (Thm. 4.3), all products $[f, g] \cdot [f, g]$ with continuous f, g in $C(K)$ with first-countable K are convex only if K is finite (Thm. 4.4).

Finally, we study square roots of intervals. A closed convex set C with $C \cdot C = [f, g]$ is called a *square root* of the interval $[f, g]$. In order that the interval $[f, g]$ have a square root which is an interval, it is necessary that $g \geq 0$ and $|f| \leq g$. If K is first-countable and if every interval $[f, g]$ with

$-g \leq f \leq 0 < g$ has a square root which is an interval, then K is finite (Thm. 5.2).

We may remark that our results, characterizing finite spaces K via properties of $C(K)$, are in line with some other results in the literature. Probably the best known assertion of this type is that K is finite if $C(K)$ has finite dimension, or that K is finite if $C(K)$ is reflexive. A less simple example is [1, Prop. 4.3.12]: Suppose that K is metrizable. If $C(K)$ is order-complete, then K is finite.

2. Preliminaries. Throughout, K is a compact Hausdorff space, and $C(K)$ is the Banach space of continuous real functions on K with the supremum norm $\|\cdot\|_\infty$. Besides the sum $f + g$ of $f, g \in C(K)$, there is also a (pointwise defined) product, denoted by fg . These operations carry over to subsets A, B of $C(K)$:

$$A + B = \{a + b : a \in A, b \in B\}, \quad A \cdot B = \{ab : a \in A, b \in B\}.$$

Recall that by an *interval* in $C(K)$ we understand a set of the form

$$[f, g] = \{h \in C(K) : f \leq h \leq g\},$$

where $f, g : K \rightarrow \mathbb{R}$ are bounded functions (generally not elements of $C(K)$) and inequalities like $f \leq h$ are defined pointwise. Every interval is bounded and, as shown in [11], closed and convex (but may be empty). To handle intervals, the notion of semicontinuity is needed.

A bounded function $f : K \rightarrow \mathbb{R}$ is *lower semicontinuous* (abbreviated by l.s.c.) if, for each $x \in K$,

$$f(x) \leq \liminf_{y \rightarrow x} f(y) := \sup_{U \in \mathcal{U}(x)} \inf_{y \in U \setminus \{x\}} f(y),$$

where $\mathcal{U}(x)$ denotes the system of neighbourhoods of x . The function f is *upper semicontinuous* (abbreviated by u.s.c.) if $-f$ is l.s.c. For a bounded function f , the function

$$f^\wedge = \sup\{h \in C(K) : h \leq f\}$$

is l.s.c. and is called the *lower semicontinuous envelope* of f . The function

$$f^\vee = \inf\{h \in C(K) : h \geq f\}$$

is u.s.c. and is called the *upper semicontinuous envelope* of f .

As shown in [11], every interval $[f, g]$ satisfies $[f, g] = [f^\vee, g^\wedge]$, and $[f^\vee, g^\wedge] \neq \emptyset$ if and only if $f^\vee \leq g^\wedge$. Thus, every nonempty interval has a *canonical representation*, that is, one of the form $[f, g]$ with f u.s.c., g l.s.c. and $f \leq g$. For the following result, proved in [11], the canonical representation is crucial.

PROPOSITION 2.1. *If $[f, g]$ and $[\varphi, \psi]$ are intervals in $C(K)$ in canonical representation, then*

$$(2.1) \quad [f, g] + [\varphi, \psi] = [f + \varphi, g + \psi].$$

For intervals in $C(K)$, the properties collected in the following definition will play an important role.

DEFINITION 2.2. An interval is called

- (i) *signed* if it has a representation $[f, g]$ where either $f > 0$ or $g < 0$,
- (ii) *positive [nonnegative]* if it has a representation $[f, g]$ with $f > 0$ [with $f \geq 0$],
- (iii) *continuous* if it has a representation $[f, g]$ with f, g continuous.

We note that for a signed interval, its canonical representation $[f, g]$ also has the property that either $f > 0$ or $g < 0$. Further, a representation $[f, g]$ of an interval with f, g continuous is its canonical representation. We also remark that in [11, Thm. 6.2], the continuous intervals have been characterized as being precisely the Minkowski summands of balls.

Turning to products, we note an obvious fact on products of real intervals, which will be used repeatedly. Let $I, J \subset \mathbb{R}$ be nonempty compact intervals. Their (algebraic) product is defined by

$$I \cdot J := \{ab : a \in I, b \in J\}.$$

(We call this product ‘algebraic’ to distinguish it from the Cartesian product.) Let

$$u = \min\{ab : a \in I, b \in J\}, \quad v = \max\{ab : a \in I, b \in J\}.$$

Then trivially $I \cdot J \subset [u, v]$. If $u = a_0b_0$ and $v = a_1b_1$ with $a_0, a_1 \in I$ and $b_0, b_1 \in J$, define, for $t \in [0, 1]$,

$$f(t) := ((1-t)a_0 + ta_1) \cdot ((1-t)b_0 + tb_1).$$

Then $f(0) = u$ and $f(1) = v$. Let $z \in [u, v]$. By continuity, there exists $\tau \in [0, 1]$ with $f(\tau) = z$, hence

$$z = ((1-\tau)a_0 + \tau a_1) \cdot ((1-\tau)b_0 + \tau b_1),$$

and here $(1-\tau)a_0 + \tau a_1 \in I$ and $(1-\tau)b_0 + \tau b_1 \in J$. It follows that $I \cdot J = [u, v]$.

3. Products of intervals in $C(K)$. Recall that K is a compact Hausdorff space and $C(K)$ is the Banach algebra of continuous real functions on K with the supremum norm. In view of the result (2.1) on the sum of two intervals, it is natural to ask about their product,

$$[f, g] \cdot [\varphi, \psi] = \{h\eta : h \in [f, g], \eta \in [\varphi, \psi]\}.$$

A particular case is the product of a scalar by an interval (the scalar can be considered as a constant function). Clearly, for an arbitrary interval $[f, g] \subset C(K)$, we have $\lambda[f, g] = [\lambda f, \lambda g]$ if $\lambda > 0$ and $\lambda[f, g] = [\lambda g, \lambda f]$ if $\lambda < 0$. The product of a scalar by the Minkowski sum of two intervals is distributive. By contrast, the product of an interval by a sum of scalars satisfies the distributive law only when the scalars have the same sign. Consider, for instance, the unit ball $[-\bar{1}, \bar{1}]$ of $C(K)$ (we denote by \bar{c} the constant function with value c). Then

$$[-\bar{1}, \bar{1}] = (-1+2)[- \bar{1}, \bar{1}] \neq (-1)[- \bar{1}, \bar{1}] + 2[- \bar{1}, \bar{1}] = [-\bar{1}, \bar{1}] + [-\bar{2}, \bar{2}] = [-\bar{3}, \bar{3}].$$

Suppose now that $[f, g]$ and $[\varphi, \psi]$ are intervals in $C(K)$. For each $x \in K$ we have

$$[f(x), g(x)] \cdot [\varphi(x), \psi(x)] = [u(x), v(x)]$$

with

$$\begin{aligned} u(x) &= \min\{ab : a \in [f(x), g(x)], b \in [\varphi(x), \psi(x)]\}, \\ v(x) &= \max\{ab : a \in [f(x), g(x)], b \in [\varphi(x), \psi(x)]\}. \end{aligned}$$

This defines bounded functions u, v on K . Trivially, we have

$$[f, g] \cdot [\varphi, \psi] \subset [u, v].$$

We call (u, v) the *pair of bounding functions* corresponding to the quadruple (f, g, φ, ψ) (although this correspondence is not shown in the notation).

LEMMA 3.1. *Let $[f, g]$ and $[\varphi, \psi]$ be intervals in canonical representation in $C(K)$, and let (u, v) be the pair of bounding functions corresponding to (f, g, φ, ψ) .*

- (i) *If $[f, g] \cdot [\varphi, \psi]$ is an interval $[F, G]$ in canonical representation and $x \in K$, then*

$$[F(x), G(x)] = [f(x), g(x)] \cdot [\varphi(x), \psi(x)].$$

- (ii) *If $[f, g] \cdot [\varphi, \psi]$ is an interval, then it is the interval $[u^\vee, v^\wedge]$.*
- (iii) *If f, g, φ, ψ are continuous, then u, v are continuous.*

Proof. First we remark that for an interval $[f, g]$ in canonical representation and for $x \in K$ and $\lambda \in \mathbb{R}$ with $f(x) \leq \lambda \leq g(x)$, there exists $h \in [f, g]$ with $h(x) = \lambda$ (see [11, Lemma 4.2]). Assertion (i) follows immediately from this remark.

To prove (ii), assume that $P := [f, g] \cdot [\varphi, \psi]$ is an interval. It is nonempty and hence has a unique canonical representation $P = [F, G]$. For each $H \in P$ we have $u \leq H \leq v$ and hence also

$$u^\vee \leq H \leq v^\wedge.$$

Since $[F, G]$ is the canonical representation, we have $F = \inf [F, G]$ and $G = \sup [F, G]$ (see [11, p. 357]) and hence $u^\vee \leq F$ and $G \leq v^\wedge$. Since $u(x)$ is an

attained minimum, there are numbers a, b with $f(x) \leq a \leq g(x)$, $\varphi(x) \leq b \leq \psi(x)$ and $ab = u(x)$. By the initial remark, there are $h \in [f, g]$ and $\eta \in [\varphi, \psi]$ such that $h(x) = a$ and $\eta(x) = b$. Then $h\eta \in P = [F, G]$ and $(h\eta)(x) = ab = u(x)$. It follows that $u(x) \geq F(x)$. We conclude that $u \geq F$, hence $u^\vee \geq F^\vee = F$ and thus $u^\vee = F$. Similarly we obtain $v^\wedge = G$. This proves (ii).

To prove (iii), let f, g, φ, ψ be continuous. The mapping

$$x \mapsto [f(x), g(x)] \times [\varphi(x), \psi(x)]$$

from K into the set of rectangles in \mathbb{R}^2 is continuous with respect to the Hausdorff metric on the space of nonempty compact subsets of \mathbb{R}^2 . The product mapping $(a, b) \mapsto ab$ from \mathbb{R}^2 to \mathbb{R} is continuous. Therefore, also $x \mapsto \min[f(x), g(x)] \cdot [\varphi(x), \psi(x)]$ is continuous, and similarly for \max . ■

We return to the question whether there is a counterpart of (2.1) for the algebraic product. It turns out that this is the case only under very strong additional assumptions. First we prove the following.

THEOREM 3.2. *The product of two signed intervals in $C(K)$ is always a signed interval.*

Proof. Let $[f, g]$ and $[\varphi, \psi]$ be intervals in $C(K)$, without loss of generality in canonical representation. We shall prove the following:

$$(3.1) \quad \text{if } f > 0, \varphi > 0, \quad \text{then } [f, g] \cdot [\varphi, \psi] = [f\varphi, g\psi],$$

$$(3.2) \quad \text{if } f > 0, \psi < 0, \quad \text{then } [f, g] \cdot [\varphi, \psi] = [g\varphi, f\psi],$$

$$(3.3) \quad \text{if } g < 0, \psi < 0, \quad \text{then } [f, g] \cdot [\varphi, \psi] = [g\psi, f\varphi],$$

$$(3.4) \quad \text{if } g < 0, \varphi > 0, \quad \text{then } [f, g] \cdot [\varphi, \psi] = [f\psi, g\varphi].$$

Suppose first that $f, \varphi > 0$. The relation $[f, g] \cdot [\varphi, \psi] \subset [f\varphi, g\psi]$ is trivial. To prove the reverse inclusion, assume that $H \in [f\varphi, g\psi]$. Let

$$\lambda = \max\{f, H/\psi\}, \quad \mu = \min\{g, H/\varphi\}.$$

Since ψ is l.s.c. and H is continuous, the function H/ψ is u.s.c. Since also f is u.s.c., the function λ is u.s.c. Similarly, μ is l.s.c. Moreover, $f \leq g$, $f \leq H/\varphi$, $H/\psi \leq g$, $H/\psi \leq H/\varphi$, hence $\lambda \leq \mu$. By the Katětov–Tong insertion theorem ([8], [12], or [7, p. 61]), there exists $h \in [\lambda, \mu]$. Then $f \leq h \leq g$, and the function $\eta := H/h$ (which can be defined since $f > 0$ implies $h > 0$) satisfies $\varphi \leq \eta \leq \psi$, thus

$$H = h\eta \in [f, g] \cdot [\varphi, \psi].$$

This completes the proof of (3.1).

Relations (3.2)–(3.4) follow from (3.1). For instance, in the case of (3.2) we have

$$[f, g] \cdot [\varphi, \psi] = -[f, g] \cdot [-\psi, -\varphi] = -[-f\psi, -g\varphi] = [g\varphi, f\psi].$$

Cases (3.3), (3.4) are settled similarly. ■

4. Properties of products of intervals. One might be inclined to think that, for example, in (3.1) the positive intervals could be replaced by nonnegative intervals. This, however, is in general impossible. Of course, if K is discrete (and thus finite, since it is compact), then for any intervals $[f, g], [\varphi, \psi]$ we have $[f, g] \cdot [\varphi, \psi] = [u, v]$ (with u, v as in Lemma 3.1), since any $H \in [u, v]$ can be written, for each $x \in K$, in the form $H(x) = h(x)\eta(x)$ with $h(x) \in [f(x), g(x)]$ and $\eta(x) \in [\varphi(x), \psi(x)]$, and every real function on K is continuous. Under a certain restriction on K , we can reverse this observation. It turns out that none of the inequalities $f > 0, \varphi > 0$ can be relaxed in general, even if f, g, φ, ψ are continuous, and even if we do not require the full interval property of the product, but only closedness.

THEOREM 4.1. *Assume that K is first-countable. If in $C(K)$ the product of a positive and a nonnegative interval, both continuous, is always closed, then K is finite.*

Proof. Assume that the assumptions are satisfied, but K is not finite. Then (since K is compact) there is an $x_0 \in K$ that is not an isolated point, that is, every neighbourhood of x_0 contains a point different from x_0 . Since K is a first-countable Hausdorff space, there is in K a decreasing sequence $(U_n)_{n \in \mathbb{N}}$ of open sets forming a neighbourhood basis of x_0 , and also a sequence $(y_n)_{n \in \mathbb{N}}$ of points satisfying $y_n \in U_n \setminus U_{n+1}$ for $n \in \mathbb{N}$. Then the sequence $(y_n)_{n \in \mathbb{N}}$ is injective and converges to x_0 .

By induction, we show the existence of a sequence $(p_n)_{n \in \mathbb{N}}$ of continuous functions $p_n : K \rightarrow [1, 2]$ with special properties. First, we set $p_1 \equiv 1$. If for the number $n \in \mathbb{N}$ the continuous function $p_n : K \rightarrow [1, 2]$ has already been defined, we use the Tietze extension theorem to find a continuous function $p_{n+1} : K \rightarrow [1, 2]$ with

$$p_{n+1}(y_{n+1}) = \begin{cases} 1 & \text{if } n+1 \text{ is odd,} \\ 2 & \text{if } n+1 \text{ is even,} \end{cases} \quad p_{n+1}(x) = p_n(x) \quad \text{for } x \in K \setminus U_{n+1}.$$

After this inductive process, for each $x \in K \setminus \{x_0\}$ there is some $m \in \mathbb{N}$ such that $x \notin U_m$ and therefore $p_k(x) = p_m(x)$ for $k \geq m$. We define $p(x) = p_m(x)$ and set $p(x_0) = 0$. Summarizing, we have:

The function $p : K \rightarrow [1, 2]$ is continuous on $K \setminus \{x_0\}$, and it satisfies

$$p(y_n) = \begin{cases} 1 & \text{if } n \text{ is odd,} \\ 2 & \text{if } n \text{ is even.} \end{cases}$$

Next, we define

$$\varphi(x_0) = 0, \quad \varphi(y_n) = 1/n \quad \text{for } n \in \mathbb{N},$$

and extend this to a continuous function $\varphi : K \rightarrow [0, 1]$. Further, we define $f, g : K \rightarrow \mathbb{R}$ by $f = \bar{1}$ and $g = \bar{2}$.

The function

$$H = p\varphi$$

is continuous on K , since p is continuous on $K \setminus \{x_0\}$, φ is continuous on K , the function p is bounded, and $\varphi(x_0) = 0$.

We claim that H is in the closure of $[f, g] \cdot [\varphi, \varphi]$. To see this, let $\varepsilon > 0$. We choose $m \in \mathbb{N}$ with $\varphi(x) \leq \varepsilon$ for $x \in U_m$. Since $p_m \in [f, g]$, the function $H_m = p_m\varphi$ satisfies $H_m \in [f, g] \cdot [\varphi, \varphi]$. Further, $H_m(x) = H(x)$ for $x \in K \setminus U_m$, while for $x \in U_m$ we get

$$|H_m(x) - H(x)| = |p_m(x) - p(x)|\varphi(x) \leq \varepsilon.$$

Thus, $\|H_m - H\|_\infty \leq \varepsilon$, so H is in the closure of $[f, g] \cdot [\varphi, \varphi]$.

Since by assumption the latter set is closed, we have $H \in [f, g] \cdot [\varphi, \varphi]$, hence there is an $h \in [f, g]$ with $H = h\varphi$. This yields $h(x) = p(x)$ whenever $\varphi(x) \neq 0$, in particular $h(y_n) = 1$ for odd n and $h(y_n) = 2$ for even n , which contradicts the continuity of h at x_0 . ■

In Theorem 4.1 (and in later theorems), first countability of K is assumed in order to ensure that for every $A \subset K$ and every $x \in \overline{A}$, there exists a sequence in A that converges to x . That some assumption of this kind cannot be avoided can be seen from the following.

Let \mathbb{N} denote the discrete space of natural numbers (it could be replaced by any infinite discrete space), and let $\beta\mathbb{N}$ be its Stone–Čech compactification (see, e.g., [6]). For $K = \beta\mathbb{N}$, the proof of Theorem 4.1 breaks down, since $\beta\mathbb{N}$ does not contain a nontrivial convergent sequence (see, e.g., [7, Corollary 3.6.15]). And indeed, in $C(\beta\mathbb{N})$ the product of two intervals defined by continuous functions is always an interval. To show this, we use the fact that there is an order-preserving linear isometry $\Phi : \ell^\infty(\mathbb{N}) \rightarrow C(\beta\mathbb{N})$, where $\ell^\infty(\mathbb{N})$ is the Banach space of all bounded real functions on \mathbb{N} with the supremum norm.

PROPOSITION 4.2. *In $C(\beta\mathbb{N})$, the product of any two continuous intervals is an interval.*

Proof. The Stone–Čech compactification $\beta\mathbb{N}$ contains a dense subspace homeomorphic to \mathbb{N} , and we can assume that \mathbb{N} is already a dense subspace of $\beta\mathbb{N}$. The linear isometry Φ is then the extension of a function in $\ell^\infty(\mathbb{N})$ to a function in $C(\beta\mathbb{N})$.

Let $[f, g]$ and $[\varphi, \psi]$ be intervals in $C(\beta\mathbb{N})$, where f, g, φ, ψ are continuous, and let (u, v) be the pair of bounding functions corresponding to the quadruple (f, g, φ, ψ) . According to Lemma 3.1, the functions u, v are also continuous. We will show that

$$(4.1) \quad [f, g] \cdot [\varphi, \psi] = [u, v].$$

The inclusion $[f, g] \cdot [\varphi, \psi] \subset [u, v]$ is clear.

Let $H \in [u, v]$. For each $n \in \mathbb{N}$ we have

$$[f(n), g(n)] \cdot [\varphi(n), \psi(n)] = [u(n), v(n)],$$

by the definition of u, v . Since $H(n) \in [u(n), v(n)]$, there are $\tilde{h}(n) \in [f(n), g(n)]$ and $\tilde{\eta}(n) \in [\varphi(n), \psi(n)]$ such that

$$\tilde{h}(n)\tilde{\eta}(n) = H(n).$$

The functions $\tilde{h}, \tilde{\eta}$ thus defined on \mathbb{N} are bounded and hence elements of $\ell^\infty(\mathbb{N})$. Let $h = \Phi(\tilde{h}), \eta = \Phi(\tilde{\eta})$ be their extensions to $C(\beta\mathbb{N})$. Since on the dense subspace \mathbb{N} the continuous functions $f, g, \varphi, \psi, h, \eta$ satisfy $f \leq h \leq g, \varphi \leq \eta \leq \psi$ and $H = h\eta$, they satisfy the same relations on $\beta\mathbb{N}$. This proves that $h \in [f, g], \eta \in [\varphi, \psi], H = h\eta$ on $\beta\mathbb{N}$ and thus $H \in [f, g] \cdot [\varphi, \psi]$, which completes the proof of (4.1). ■

In contrast, in $C(\alpha\mathbb{N})$, where $\alpha\mathbb{N}$ denotes the one-point compactification of \mathbb{N} , the product of two continuous intervals need not be an interval. This follows from Theorem 4.1, since $\alpha\mathbb{N}$ is a first-countable compact Hausdorff space.

Intervals are closed and convex. As the product of two intervals need not be closed, even if they are nonnegative, it is a natural question whether it must at least be convex. That this is not a trivial question can be seen from Theorem 4.4 below, where the nonnegativity assumption is deleted and convexity does not hold in general. With the nonnegativity assumption, however, convexity can be proved.

THEOREM 4.3. *The product of two nonnegative intervals in $C(K)$ is always convex.*

Proof. Let $0 \leq f \leq g$ and $0 \leq \varphi \leq \psi$ be bounded functions, and let $H_1, H_2 \in [f, g] \cdot [\varphi, \psi]$. Thus there are $h_1, h_2 \in [f, g]$ and $\eta_1, \eta_2 \in [\varphi, \psi]$ such that $H_i = h_i\eta_i$ for $i = 1, 2$. Let $0 < t < 1$. We have to show that

$$H := (1-t)H_1 + tH_2 = (1-t)h_1\eta_1 + th_2\eta_2$$

satisfies

$$(4.2) \quad H \in [f, g] \cdot [\varphi, \psi].$$

To prepare this, let $M = \{(a, \alpha, b, \beta) \in \mathbb{R}^4 : a, \alpha, b, \beta \geq 0\}$ and $z = (a, \alpha, b, \beta) \in M$. We consider (a, α) and (b, β) as points in \mathbb{R}^2 with the standard basis. They are opposite vertices of a unique rectangle (possibly degenerate) with edges parallel to the coordinate axes. We denote this rectangle by $\text{Rec}(z)$. Further, we define the hyperbola

$$\text{Hyp}(z) = \{(c, \gamma) \in \mathbb{R}^2 : c\gamma = (1-t)a\alpha + tb\beta\}$$

(which may be degenerate). The function p defined by

$$p(\tau) = ((1-\tau)a + \tau b) \cdot ((1-\tau)\alpha + \tau\beta), \quad 0 \leq \tau \leq 1,$$

satisfies $p(0) = a\alpha$ and $p(1) = b\beta$, hence there is some $\tau_0 \in [0, 1]$ with $p(\tau_0) = (1-t)a\alpha + t b\beta$. Thus, $\text{Rec}(z)$ has a nonempty intersection with $\text{Hyp}(z)$. We will find a continuous map $P : M \rightarrow \mathbb{R}^2$ satisfying, for all $z \in M$,

$$P(z) \in \text{Rec}(z) \cap \text{Hyp}(z).$$

Having such a map, we can proceed as follows. For $x \in K$, we define

$$z(x) = (h_1(x), \eta_1(x), h_2(x), \eta_2(x)).$$

Then we define $\mathcal{P}(x) = P(z(x))$ for $x \in K$. The mapping $\mathcal{P} : K \rightarrow \mathbb{R}^2$ is continuous and satisfies $\mathcal{P}(x) \in \text{Rec}(z(x)) \cap \text{Hyp}(z(x))$. Hence, if we define functions h, η on K by $\mathcal{P}(x) = (h(x), \eta(x))$, then h and η are continuous, they satisfy $h(x)\eta(x) = H(x)$ since $\mathcal{P}(x) \in \text{Hyp}(z(x))$, and $f \leq h \leq g$, $\varphi \leq \eta \leq \psi$ since $\mathcal{P}(x) \in \text{Rec}(z(x))$ and the latter rectangle is contained in $\text{Rec}(f(x), g(x), \varphi(x), \psi(x))$. Thus, $h \in [f, g]$, $\eta \in [\varphi, \psi]$ and $h\eta = H$, which shows that $H \in [f, g] \cdot [\varphi, \psi]$, proving (4.2).

We proceed to construct the mapping P . First, if $a = b = 0$, we define

$$P(z) = (0, (1-t)\alpha + t\beta).$$

If $\alpha = \beta = 0$, we set

$$P(z) = ((1-t)a + t b, 0).$$

Now we assume that $\max\{a, b\} > 0$ and $\max\{\alpha, \beta\} > 0$. Then $(1-t)a + t b \neq 0$ and $(1-t)\alpha + t\beta \neq 0$, hence we can define

$$(4.3) \quad A = \left(\frac{(1-t)a\alpha + t b\beta}{(1-t)\alpha + t\beta}, (1-t)\alpha + t\beta \right),$$

$$(4.4) \quad B = \left((1-t)a + t b, \frac{(1-t)a\alpha + t b\beta}{(1-t)a + t b} \right).$$

The point A belongs to $\text{Hyp}(z)$ (trivially) and to $\text{Rec}(z)$ (observe that its first coordinate is a convex combination of a and b , and its second coordinate is a convex combination of α and β). Similarly, $B \in \text{Hyp}(z) \cap \text{Rec}(z)$. The point

$$(4.5) \quad C = \frac{((1-t)\alpha + t\beta)A + ((1-t)a + t b)B}{(1-t)\alpha + t\beta + (1-t)a + t b},$$

is a convex combination of A and B and hence belongs to $\text{Rec}(z)$. Let L be the line in \mathbb{R}^2 that is parallel to the vector $(1, 1)$ and passes through C . Then either the points A and B coincide, then $C = A = B$ and hence C lies on $\text{Hyp}(z)$, or A and B lie on different sides of L . Since the part of $\text{Hyp}(z)$ between A and B lies in $\text{Rec}(z)$, the line L intersects $\text{Hyp}(z)$ in a unique point belonging to $\text{Rec}(z)$; we denote this point by $P(z)$.

We have to show that the mapping P thus defined is continuous. Let $z = (a, \alpha, b, \beta) \in M$. Suppose first that $\max\{a, b\} > 0$ and $\max\{\alpha, \beta\} > 0$.

Then there is a neighbourhood V of z in M such that $\max\{a', b'\} > 0$ and $\max\{\alpha', \beta'\} > 0$ for all $z' = (a', \alpha', b', \beta') \in V$. Since A, B, C defined above are continuous functions of a, α, b, β , and also the intersection point of L and Hyp defines a continuous function, the mapping P is continuous at z .

Second, suppose that $z = (0, 0, 0, 0)$. Then $\text{Rec}(z) = \{(0, 0)\}$ and $P(z) = (0, 0)$. Let U be a neighbourhood of $(0, 0)$ in \mathbb{R}^2 . There is a neighbourhood V of z in M such that for all $z' \in M$ we have $\text{Rec}(z') \subset U$ and hence $P(z') \in U$. Thus, P is continuous at z .

Third, let $z = (a, \alpha, b, \beta) \in M$ belong to one of the remaining cases, that is, either $\alpha = \beta = 0$, $\max\{a, b\} > 0$, or $a = b = 0$, $\max\{\alpha, \beta\} > 0$. Suppose the former holds. By definition, $P(z) = ((1-t)a + tb, 0)$. Let U be a neighbourhood of $P(z)$ in \mathbb{R}^2 . Let $z_i = (a_i, \alpha_i, b_i, \beta_i) \in M$, $i \in \mathbb{N}$, be such that $z_i \rightarrow z$ for $i \rightarrow \infty$. We may assume that $\max\{a_i, b_i\} > 0$, since this holds for all sufficiently large i . If $\alpha_i = \beta_i = 0$, then $P(z_i) = ((1-t)a_i + tb_i, 0)$, which is in U for sufficiently large i . In the following, we consider the indices i for which $(\alpha_i, \beta_i) \neq (0, 0)$. First we notice that the parameter $(1-t)a_i\alpha_i + tb_i\beta_i$ of the hyperbola $\text{Hyp}(z_i)$ tends to zero as $i \rightarrow \infty$. Therefore, there is a neighbourhood $U' \subset U$ with the following property. If $C \in U'$, then the line in \mathbb{R}^2 that is parallel to the vector $(1, 1)$ and passes through C intersects $\text{Hyp}(z_i)$ inside U . Define A_i, B_i, C_i by (4.3), (4.4), (4.5) with z replaced by z_i . Then $B_i \rightarrow ((1-t)a + tb, 0)$, and A_i remains bounded, since f, g, φ, ψ are bounded. The weight of A_i in

$$C_i = \frac{((1-t)\alpha_i + t\beta_i)A_i + ((1-t)a_i + tb_i)B_i}{(1-t)\alpha_i + t\beta_i + (1-t)a_i + tb_i}$$

tends to zero as $i \rightarrow \infty$, and so $C_i \rightarrow ((1-t)a + tb, 0)$. Therefore, $C_i \in U'$ and hence $P(z_i) \in U$ for sufficiently large i . This shows that P is continuous at z . The case $a = b = 0$, $\max\{\alpha, \beta\} > 0$ is treated similarly. ■

In the proof of Theorem 4.1, it was important that we can take products of different intervals. In fact, $[f, g] \cdot [f, g]$ with $f \geq 0$ is always an interval, since for $H \in [f^2, g^2]$, the function $h = H^{1/2}$ satisfies $h \in [f, g]$ and $h^2 = H$. That the assumption $f \geq 0$ cannot be deleted is shown by the following theorem. It reveals that in general the product $[f, g] \cdot [f, g]$, even for f, g continuous, need not be an interval, because it need not even be convex.

THEOREM 4.4. *Assume that K is first-countable. If for any continuous interval $[f, g]$ in $C(K)$ the product $[f, g] \cdot [f, g]$ is convex, then K is finite.*

Proof. Assume that the assumptions are satisfied, but K is not finite. Then there is an $x_0 \in K$ which is not an isolated point. Since K is a first-countable Hausdorff space, there is an injective sequence $(y_n)_{n \in \mathbb{N}}$ in $K \setminus \{x_0\}$ that converges to x_0 . The set $X = \{y_n : n \in \mathbb{N}\} \cup \{x_0\}$ is closed. Define

$f : X \rightarrow [-2, 0]$ and $g : X \rightarrow [0, 2]$ by

$$f(y_n) = -1 + (-1)^n/n, \quad g(y_n) = 1 + (-1)^n/n \quad \text{for } n \in \mathbb{N}$$

and $f(x_0) = -1, g(x_0) = 1$. Since K is a normal space, there exist continuous extensions $f : K \rightarrow [-2, 0]$ and $g : K \rightarrow [0, 2]$. Then $f \leq g$ on K .

If we define

$$H = \frac{1}{2}f^2 + \frac{1}{2}g^2,$$

then

$$H(y_n) = 1 + \frac{1}{n^2}$$

for every $n \in \mathbb{N}$. Since $f^2, g^2 \in [f, g] \cdot [f, g]$ and the latter set is convex by assumption, we have

$$H \in [f, g] \cdot [f, g].$$

Therefore, there are $h, \eta \in [f, g]$ such that $h\eta = H$. For each $n \in \mathbb{N}$ we have

$$h(y_n)\eta(y_n) = 1 + \frac{1}{n^2}.$$

In particular, $h(y_n)$ and $\eta(y_n)$ have the same sign.

Let n be even. If $h(y_n)$ and $\eta(y_n)$ are both negative, then

$$h(y_n)\eta(y_n) \leq \left(-1 + \frac{1}{n}\right)^2 < 1 + \frac{1}{n^2},$$

a contradiction. Therefore, $h(y_n)$ and $\eta(y_n)$ are both positive, which implies

$$1 + \frac{1}{n^2} = h(y_n)\eta(y_n) \leq h(y_n)g(y_n) = h(y_n)\left(1 + \frac{1}{n}\right)$$

and thus $h(y_n) \geq 1 - 1/n$.

Let n be odd. If $h(y_n)$ and $\eta(y_n)$ are both positive, then

$$h(y_n)\eta(y_n) \leq \left(1 - \frac{1}{n}\right)^2 < 1 + \frac{1}{n^2},$$

a contradiction. Therefore, $h(y_n)$ and $\eta(y_n)$ are both negative, and

$$1 + \frac{1}{n^2} = |h(y_n)\eta(y_n)| \leq |h(y_n)||\eta(y_n)| = |h(y_n)|\left(1 + \frac{1}{n}\right),$$

hence $h(y_n) \leq -1 + 1/n$.

Thus, every neighbourhood of x_0 contains points where h is arbitrarily close to 1 and points where h is arbitrarily close to -1 . Since h is continuous, this is a contradiction. ■

5. Square roots of intervals. A *convex body* in $C(K)$ is a nonempty bounded closed convex subset of $C(K)$. In [11, Theorem 4.3], it was shown that if $C, D \subset C(K)$ are convex bodies such that the closure of $C + D$ is

an interval, then each of C and D is an interval. One may ask whether an analogous result is true for products instead of sums. But it is easily seen that this is not the case (see the example below).

Now we restrict ourselves to products of a convex body with itself, and we say that the convex body C is a *square root* of the interval $[f, g]$ if $C \cdot C = [f, g]$. It is easy to see that an interval can have a square root that is not an interval, provided that K is not a singleton. For example, let x_0, x_1 be distinct points of K , and let

$$A = \{f \in [-\bar{1}, \bar{1}] : f(x_0) \geq 0\}, \quad B = [-\bar{1}, -\bar{1}].$$

Then A and B are convex bodies in $C(K)$. Let C be the convex hull of $A \cup B$; it is closed. Then $C \cdot C \subset [-\bar{1}, \bar{1}]$ trivially. Conversely, $f \in [-\bar{1}, \bar{1}]$ satisfies either $f(x_0) \geq 0$ and hence $f = f \cdot \bar{1} \in C \cdot C$, or $f(x_0) < 0$ and hence $f = (-f) \cdot (-\bar{1}) \in C \cdot C$. Thus, C is a square root of $[-\bar{1}, \bar{1}]$. On the other hand, C is not an interval: if it were, it could only be $[-\bar{1}, \bar{1}]$, but a function $h \in [-\bar{1}, \bar{1}]$ with $h(x_0) = -1$ and $h(x_1) = 1$ (which exists by Urysohn's lemma) does not belong to C .

This raises the question: which intervals have a square root that is an interval? Trivially, this holds for $[f, g]$ with $f \geq 0$. In fact, if $f \geq 0$ and if $H \in [f, g]$, then $\sqrt{H} \in [\sqrt{f}, \sqrt{g}]$, which shows that $[\sqrt{f}, \sqrt{g}] \cdot [\sqrt{f}, \sqrt{g}] = [f, g]$.

The following result provides additional information, which will be used below to show that also assuming the existence of square roots leads to finite spaces.

PROPOSITION 5.1. *Let $[f, g]$ be an interval in canonical representation in $C(K)$. In order that $[f, g]$ have a square root which is an interval, it is necessary that*

$$(5.1) \quad g \geq 0 \quad \text{and} \quad |f| \leq g,$$

and it is sufficient that g is continuous and

$$(5.2) \quad -g \leq f \leq 0 < g.$$

Proof. To prove the necessity, consider first an interval $[a, b]$ with $a < b$ in \mathbb{R} , and suppose that it has a square root (that is, an interval $[c, d]$ with $[c, d] \cdot [c, d] = [a, b]$). Then $b \geq 0$ trivially. If $a \geq 0$, then $[\sqrt{a}, \sqrt{b}]$ and $[-\sqrt{b}, -\sqrt{a}]$ are the square roots of $[a, b]$. Let $a < 0 \leq b$. A square root of $[a, b]$ must be of the form $[c, d]$ with $c \leq 0 \leq d$. If $d \geq |c|$, then $[c, d] \cdot [c, d] = [cd, d^2]$, hence $[a/\sqrt{b}, \sqrt{b}]$ is a square root of $[a, b]$. If $d < |c|$, then $[c, d] \cdot [c, d] = [cd, c^2]$, hence $[-\sqrt{b}, -a/\sqrt{b}]$ is a square root of $[a, b]$. In either case, $[a/\sqrt{b}, \sqrt{b}]$ and $[-\sqrt{b}, -a/\sqrt{b}]$ are square roots of $[a, b]$, and we have $|a| \leq b$.

Now let $[f, g]$ be an interval in canonical representation in $C(K)$. Suppose it has a square root which is an interval. Then we can assume that the latter

is also in canonical representation. Now it follows from Lemma 3.1(1), by applying the preceding observation to $[a, b] = [f(x), g(x)]$ for $x \in K$, that condition (5.1) is necessary, as stated.

To prove the sufficiency, suppose that (5.2) is satisfied and that g is continuous. Define, for $x \in K$,

$$\varphi(x) = f(x)/\sqrt{g(x)}, \quad \psi(x) = \sqrt{g(x)}.$$

For each $x \in K$ we have $[\varphi(x), \psi(x)] \cdot [\varphi(x), \psi(x)] = [f(x), g(x)]$. This shows that $[\varphi, \psi] \cdot [\varphi, \psi] \subset [f, g]$. If $H \in [f, g]$, the functions $h = \sqrt{g}$ and $\eta = H/\sqrt{g}$ satisfy $h, \eta \in [\varphi, \psi]$ and $h\eta = H$. It follows that $[\varphi, \psi] \cdot [\varphi, \psi] = [f, g]$. ■

The continuity in the second part of Proposition 5.1 is essential, as the following theorem shows.

THEOREM 5.2. *Let K be first-countable. If every interval $[f, g]$ in canonical representation in $C(K)$ satisfying (5.2) has a square root which is an interval, then K is finite.*

Proof. Let K be first-countable, and suppose every interval in canonical representation satisfying (5.2) has a square root which is an interval. Assume that K is not finite. Then K has a point x_0 that is not isolated. Let $(y_n)_{n \in \mathbb{N}}$ be an injective sequence in $K \setminus \{x_0\}$ converging to x_0 . We define $f = -1/2$ on K and

$$g(x) = \begin{cases} 1/2 & \text{for } x = y_n \text{ with odd } n \text{ and for } x = x_0, \\ 1 & \text{for the remaining } x \in K. \end{cases}$$

Then g is l.s.c., and $[f, g]$ is an interval in canonical representation satisfying (5.2). By assumption, it has a square root which is an interval, say $[\varphi, \psi]$, without loss of generality also in canonical representation. Therefore, in particular, the function f has a representation $f = h\eta$ with $h, \eta \in [\varphi, \psi]$.

Let $x = y_n$ with n even. Then $[f(x), g(x)] = [-1/2, 1]$. As shown in the proof of Proposition 5.1, either $[\varphi(x), \psi(x)] = [-1/2, 1]$ or $[\varphi(x), \psi(x)] = [-1, 1/2]$. Since $-1/2 = f(x) = h(x)\eta(x)$, this yields

$$h(x) \in \{\pm 1/2, \pm 1\}.$$

Let $x = y_n$ with n odd. Then $[f(x), g(x)] = [-1/2, 1/2]$ and $[\varphi(x), \psi(x)] = [-1/\sqrt{2}, 1/\sqrt{2}]$. Since $-1/2 = f(x) = h(x)\eta(x)$, this yields

$$h(x) \in \{\pm 1/\sqrt{2}\}.$$

Thus, in every neighbourhood of x_0 , there are points (y_n with sufficiently large even n) where h attains one of the values $\pm 1/2, \pm 1$, and points (y_n with sufficiently large odd n) where h attains one of the values $\pm 1/\sqrt{2}$. This contradicts the continuity of h . ■

We remark that, in analogy to Proposition 4.2, in $C(\beta\mathbb{N})$ any interval $[f, g]$ with f, g continuous satisfying (5.2) has a square root which is an interval.

Acknowledgements. This research was partly supported by Ministerio de Ciencia e Innovación, grant MTM2012-34341.

We thank the anonymous referee for a thorough report, and we highly appreciate his/her valuable suggestions for improvements.

References

- [1] F. Albiac and N. J. Kalton, *Topics in Banach Space Theory*, Grad. Texts in Math. 233, Springer, New York, 2006.
- [2] M. Balcerzak, W. Wachowicz and W. Wilczyński, *Multiplying balls in the space of continuous functions on $[0, 1]$* , Studia Math. 170 (2005), 203–209.
- [3] E. Behrends, *Walk the dog, or: products of open balls in the space of continuous functions*, Funct. Approx. Comment. Math. 44 (2011), 153–164.
- [4] E. Behrends, *Products of n open subsets in the space of continuous functions on $[0, 1]$* , Studia Math. 204 (2011), 73–95.
- [5] E. Behrends, *Where is pointwise multiplication in real CK -spaces locally open?*, Fund. Math., to appear.
- [6] N. L. Carothers, *A Short Course on Banach Space Theory*, Cambridge Univ. Press, Cambridge, 2005.
- [7] R. Engelking, *General Topology*, Heldermann, Berlin, 1989.
- [8] M. Katětov, *On real-valued functions in topological spaces*, Fund. Math. 38 (1951), 85–91; Correction, Fund. Math. 40 (1953), 203–205.
- [9] A. Komisarski, *A connection between multiplication in $C(X)$ and the dimension of X* , Fund. Math. 189 (2006), 149–154.
- [10] S. Kowalczyk, *Weak openness of multiplication in the space $C(0, 1)$* , Real Anal. Exchange 35 (2009/2010), 235–242.
- [11] J. P. Moreno and R. Schneider, *Some geometry of convex bodies in $C(K)$ spaces*, J. Math. Pures Appl. 103 (2015), 352–373.
- [12] H. Tong, *Some characterizations of normal and perfectly normal spaces*, Duke Math. J. 19 (1952), 289–292.

José Pedro Moreno
Departamento de Matemáticas
Facultad de Ciencias
Universidad Autónoma de Madrid
E-28049 Madrid, Spain
E-mail: josepedro.moreno@uam.es

Rolf Schneider
Mathematisches Institut
Albert-Ludwigs-Universität
D-79104 Freiburg i. Br., Germany
E-mail: rolf.schneider@math.uni-freiburg.de