

UNTHREADABILITY WITH SMALL CONDITIONS

MAXWELL LEVINE

ABSTRACT. We introduce a forcing that adds a $\square(\aleph_2, \aleph_0)$ -sequence with countable conditions under CH. Assuming the consistency of a weakly compact cardinal, we can find a forcing extension by our new poset in which both $\square(\aleph_2, < \aleph_0)$ and $\square_{\aleph_1, \aleph_0}$ fail in the forcing extension.

1. INTRODUCTION

The tension between the stationary reflection principles of large cardinals and the fine-structural combinatorics that hold in canonical inner models is a prominent theme in set theory. In particular, variations of Jensen's principle \square_κ , which holds for all cardinals κ in Gödel's Constructible Universe L , have been studied widely. The construction of new models realizing some compatibility between large cardinal properties and square properties can help us develop a more vivid picture of the individual cardinals and their relationships to one another.

The principle \square_κ and its relatives are known as *square principles*. Each such principle asserts that there exists a coherent sequence $\mathcal{C} = \langle \mathcal{C}_\alpha : \alpha < \lambda \rangle$ where \mathcal{C}_α is a set of closed unbounded sets in α , and that \mathcal{C} does not have a *thread*, meaning that there is no club $D \subseteq \lambda$ such that $D \cap \alpha \in \mathcal{C}_\alpha$ for all limit points $\alpha \in D$. Square principles of the form $\square_{\mu, \kappa}$ assert the existence of such a \mathcal{C} of length μ^+ (notice the discrepancy in notation) in which the members of the \mathcal{C}_α 's have order-type bounded by μ , which serves as the reason why there cannot be a thread. On the other hand, square principles of the form $\square(\mu^+, \kappa)$ assert the existence of such a \mathcal{C} of length μ^+ where the non-existence of the thread is simply declared outright. The consequences of this distinction are significant for this paper.

We will follow a strain of research having to do with adding square sequences of length μ^+ with forcings consisting of conditions of cardinality $< \mu$. The poset introduced here adds a $\square(\aleph_2, \aleph_0)$ -sequence with countable conditions. This is different from the conventional way of forcing $\square(\mu^+, \kappa)$ -sequences, which uses conditions of cardinality μ . This approach was introduced by Jensen and studied thoroughly by Cummings, Foreman, and Magidor. We will refer to this as *Jensen's method* for forcing square sequences, since we will have reason to compare and contrast it with the approach used here, especially regarding the behavior of the complementary threading forcing.

The idea of the forcing described in this paper is roughly based on Baumgartner's forcing to add a club to \aleph_1 with finite sequences, except that we

use its generalization to \aleph_2 , and we crucially use a presentation that is due to Abraham [1]. More precisely, one of the main hurdles for adding a new object of size \aleph_2 with countable conditions is finding a way to preserve \aleph_2 knowing that that new \aleph_1 -sequences will be added. In our case we use elementary submodels to guarantee preservation of \aleph_2 . This use of elementary submodels is alluded to by Mitchell when he discusses the options for adding a new closed unbounded set to \aleph_2 in terms of possibilities for the sizes of conditions [9]. It also distinguishes the forcing presented in this paper from the Shelah-Stanley poset for adding \square_{\aleph_1} with countable conditions, which uses a chain condition to preserve the cardinal that is turned in to \aleph_2 [12]. (The Shelah-Stanley poset adds a strictly stronger square sequence as well.) The technical difficulty comes mostly in showing that the new forcing is countably closed, or to be more precise, that it has a dense countably closed subset. This property also contrasts the new forcing with the forcings for adding \square_{\aleph_1} using finite conditions originally used by Dolinar and Džamonja [4] and later streamlined by Neeman [11] since those cannot be countably closed. Above all, the motivation for the work here is to provide a new and distinct model in which a much-studied combinatorial principle holds.

Theorem 1. *If $V \models \text{CH}$ then there is a forcing \mathbb{S} consisting of countable conditions such that $V[\mathbb{S}] \models \square(\aleph_2, \aleph_0)$. Moreover, if $W \models$ “ κ is weakly compact”, then $W[\text{Col}(\aleph_1, < \kappa)][\mathbb{S}] \models \neg \square(\aleph_2, < \aleph_0) \wedge \neg \square_{\aleph_1, \aleph_0}$.*

We expect that the construction presented in this paper generalizes to higher cardinals to get a $\square(\kappa^{++}, \kappa)$ -sequence under the assumption that $2^\kappa = \kappa^+$. The assumption of a weakly compact cardinal is optimal because work of Jensen and Todorćević shows that this is required to make $\square(\kappa, 1)$ fail for regular κ [13].

This paper is organized as follows: The remainder of the introduction will cover background. We will introduce \mathbb{S} in the second section and show that it adds a $\square(\aleph_2, \aleph_0)$ -sequence. The third section will introduce the new threading forcing \mathbb{T} and show how it can be used with large cardinals to make stronger square principles fail.

We assume familiarity with the technique of forcing [7]. As for our conventions: Given a set of ordinals X , $\text{Lim}(X)$ is the set of limit ordinals in X , $\text{lim}(X) = \{\alpha \in X : X \cap \alpha \text{ is unbounded}\}$, $\text{lim}^+(X) = \{\alpha \leq \sup(X) : X \cap \alpha \text{ is unbounded}\}$, and $j[X] = \{j(\alpha) : \alpha \in X\}$ if j is a function. We also let $\text{ot}(X)$ denote the order-type of X , and $X \cap \text{cof}(\tau) = \{\alpha \in X : \text{cf}(\alpha) = \tau\}$. If \mathbb{P} is a forcing poset and $p, q \in \mathbb{P}$, then $p \leq q$ means that p has more information than q . If we say that \mathbb{P} is κ -distributive for a regular κ , we mean that \mathbb{P} does not add new functions $f : \lambda \rightarrow \text{ON}$ for $\lambda < \kappa$, and when we say that \mathbb{P} is κ -closed, we mean that it is closed under $\leq_{\mathbb{P}}$ -decreasing sequences of length λ for all $\lambda < \kappa$. The notation $V[\mathbb{P}]$ will refer to an extension by an unspecified generic. When depicting an iteration $\mathbb{P}_0 * \dot{\mathbb{P}}_1$, we will often drop the dot and write $\mathbb{P}_0 * \mathbb{P}_1$.

1.1. Definitions of Squares. First we define square principles of the form $\square(\lambda, \kappa)$. These are examples of incompactness principles, because they imply the non-existence of an object that appears to be approximated, namely a thread.¹

Definition 2. Let κ and λ be regular cardinals such that $\kappa < \lambda$. Then $\langle \mathcal{C}_\alpha : \alpha \in \text{Lim}(\lambda) \rangle$ is a $\square(\lambda, \kappa)$ -sequence if the following hold for all $\alpha \in \text{Lim}(\lambda)$:

- (1) \mathcal{C}_α consists of clubs in α .
- (2) $1 \leq |\mathcal{C}_\alpha| \leq \kappa$.
- (3) For all $C \in \mathcal{C}_\alpha$ and all $\beta \in \text{lim}(C)$, $C \cap \beta \in \mathcal{C}_\beta$.
- (4) There is no club $D \subset \lambda$ such that for all $\alpha \in \text{lim}(D)$, $D \cap \alpha \in \mathcal{C}_\alpha$.

If there is a $\square(\lambda, \kappa)$ -sequence, then we say that $\square(\lambda, \kappa)$ holds. If $V \models \text{“}\vec{\mathcal{C}}$ is a $\square(\lambda, \kappa)$ sequence” and $W \supset V$ is a model containing some D such that $D \cap \alpha \in \mathcal{C}_\alpha$ for all $\alpha \in \text{lim}(D)$, then we say that D threads $\vec{\mathcal{C}}$.

Definition 3. Let κ and μ be cardinals. Then $\langle \mathcal{C}_\alpha : \alpha \in \text{Lim}(\mu^+) \rangle$ is a $\square_{\mu, \kappa}$ -sequence if it is a $\square(\mu^+, \kappa)$ -sequence, and moreover for all $\alpha \in \text{Lim}(\mu^+)$ and $C \in \mathcal{C}_\alpha$, $\text{ot } C \leq \mu$.

Remark 4. If \mathcal{C} is a $\square_{\mu, \kappa}$ -sequence, then there is no $D \subset \mu^+$ such that for all $\alpha \in \text{lim}(D)$, $D \cap \alpha \in \mathcal{C}_\alpha$. (Hence point (4) from the definition of $\square(\lambda, \kappa)$ -sequences is implied.) This is because if γ were the $(\kappa + \omega)^{\text{th}}$ point of D , then $D \cap \gamma \in \mathcal{C}_\gamma$ has an order-type larger than κ .

Definition 5. If \mathcal{C} is a $\square(\lambda, \kappa)$ -sequence, we say that a poset \mathbb{P} threads \mathcal{C} if \mathbb{P} forces that there is a thread of \mathcal{C} .

1.2. Elementary Submodels and Generic Conditions. In this subsection we will outline the basic arguments we use to preserve \aleph_2 . In this regard, there are a couple of important conventions we will use:

- In the context of discussing a poset \mathbb{P} , we will refer to a regular cardinal as “sufficiently large” when $H(\Theta)$ contains enough information about \mathbb{P} for the argument at hand. In this paper we mean that $H(\Theta)$ contains \mathbb{P} and all of its antichains. More generally, we will refer to a structure $K \models \text{ZFC} - \text{Powerset}$ as “sufficiently rich” when it contains \mathbb{P} and its antichains.
- Given a poset \mathbb{P} and a sufficiently rich structure K , we will refer to an elementary submodel $M \prec K$ as *basic* if $\mathbb{P} \in M$, $|M| = \aleph_1$, $M \cap \aleph_2 \in \aleph_2$, and $M^\omega \subseteq M$, i.e. M is closed under countable sequences. This usage is specific to this paper because we will use it frequently.

Proposition 6. *Suppose CH holds, $|X| = \aleph_1$, and K is a sufficiently rich structure with $X, \aleph_2 \in K$. Then there is a basic model $M \prec K$ such that $X \subset M$.*

Proof. Let $M_0 \prec K$ be an elementary submodel of cardinality \aleph_1 such that $X \subset M_0$. Then take a chain $\langle M_i : i < \aleph_1 \rangle$ of \aleph_1 -sized elementary submodels

¹See a survey of Cummings for background on square principles of the form $\square_{\mu, \kappa}$ and stationary reflection [2].

of K such that for all $i < \aleph_1$, $\sup(M_i \cap \aleph_2) \subseteq M_{i+1}$, $M_i^\omega \subseteq M_{i+1}$ (this is where we use CH), and such that there is continuity in the sense that $M_i = \bigcup_{j < i} M_j$ for all limits $i < \aleph_1$. Then let $M = \bigcup_{i < \aleph_1} M_i$. \square

Definition 7. Let \mathbb{P} be a poset, let K be a sufficiently rich structure, and let $M \prec K$ such that $\mathbb{P} \in M$.

- A condition q is (M, \mathbb{P}) -generic if for every maximal antichain $A \in M$ (such that $A \subseteq \mathbb{P}$), for all $q' \leq q$, there is some $q'' \leq q'$ such that $q'' \leq p$ for some $p \in A \cap M$. Equivalently, q is (M, \mathbb{P}) -generic if $q \Vdash \dot{G} \cap M$ is $\mathbb{P} \cap M$ -generic over M [7].
- A condition q is *strongly* (M, \mathbb{P}) -generic if for all $q' \leq q$, there is some $p \in \mathbb{P} \cap M$ such that for every $p' \leq p$ with $p' \in \mathbb{P} \cap M$, there is some $q'' \leq q'$ such that $q'' \leq p'$. Equivalently, q is strongly (M, \mathbb{P}) -generic if $q \Vdash \dot{G} \cap M$ is $\mathbb{P} \cap M$ -generic over the ground model V [10].

It is clear that all strongly (M, \mathbb{P}) -generic conditions are (M, \mathbb{P}) -generic for sufficiently rich structures. We distinguish between these two types of generic conditions for the sake of some remarks that we will make below.

Proposition 8. *Suppose \mathbb{P} is a poset, K is a sufficiently rich structure, and $M \prec K$ with $\mathbb{P} \in M$. Suppose q is (M, \mathbb{P}) -generic and G is \mathbb{P} -generic over V with $q \in G$. Then the following hold:*

- (1) $M[G] \prec K[G]$.
- (2) $V \cap M[G] = M$.
- (3) If $\dot{f} \in M$ is such that $\Vdash_{\mathbb{P}} \dot{f} : \check{\kappa}_1 \rightarrow \check{\kappa}_2$ where $\kappa_1 \subset M$ and $\kappa_2 \in M$, then $\text{range}(\dot{f}[G]) \subset M$.

Proof. These are standard arguments,² so we will prove only (3) since (1) and (2) use the same trick. Work in V and let $\alpha \in \kappa_1$. Let $A \in M$ be a maximal antichain consisting of conditions deciding $\dot{f}(\check{\alpha})$. By elementarity there is such an $A \in M$. Given some $q' \leq q$, let $q'' \leq q'$ be below some $p \in A \cap M$ using the fact that q is an (M, \mathbb{P}) -generic condition. By elementarity, $p \Vdash \dot{f}(\check{\alpha}) = \check{\beta}$ for some $\beta \in M$, so the same information is forced by q'' . \square

Definition 9. Suppose $\mathbb{P} \in M \prec K$ for K sufficiently rich. We say that M has *generic conditions for \mathbb{P}* (resp. *strongly generic conditions*) if for all $p \in \mathbb{P} \cap M$, there is some $q \leq p$ such that q is (M, \mathbb{P}) -generic (resp. strongly (M, \mathbb{P}) -generic).

Proposition 10. *Let $M \prec K$ be a basic elementary submodel of a sufficiently rich structure such that M has generic conditions for \mathbb{P} . Then \mathbb{P} preserves \aleph_2 .*

Proof. Working in V , we have $\aleph_2 \in M$ by elementarity and $\aleph_1 \subset M$ by $M^\omega \subset M$. M would contain a \mathbb{P} -name \dot{f} for a supposed collapsing function $f : \aleph_1^V \rightarrow \aleph_2^V$ by elementarity. Apply Proposition 8. \square

²See the chapter on proper forcing in Jech [7].

2. DEFINING THE MAIN POSET

The goal of this section is to introduce our poset for adding a $\square(\aleph_2, \aleph_0)$ -sequence with countable conditions.

Definition 11. Let \mathfrak{S} be the set of $(\langle \alpha, x \rangle, [\beta, \gamma])$ such that $\alpha < \aleph_2$ is a limit ordinal, $x \in \omega$, and $[\beta, \gamma]$ is a closed interval of ordinals such that $-1 \leq \beta \leq \gamma < \alpha$.³ Define \mathbb{S} to be the set of $s \subset \mathfrak{S}$ such that the following hold:

- (1) The set s is countable.
- (2) If $(\langle \alpha, x \rangle, [\beta, \gamma]), (\langle \alpha, x \rangle, [\beta', \gamma']) \in s$, then either $\beta = \beta'$ or $[\beta, \gamma] \cap [\beta', \gamma'] = \emptyset$.
- (3) If $\text{cf}(\alpha) = \omega_1$, then for all $x, y \in \omega$ and $\beta, \gamma < \alpha$, $(\langle \alpha, x \rangle, [\beta, \gamma]) \in s$ if and only if $(\langle \alpha, y \rangle, [\beta, \gamma]) \in s$.
- (4) If $\{(\langle \alpha, x \rangle, [\beta_i, \gamma_i]) : i < \omega\} \subseteq s$ and $\beta^* = \sup_{i < \omega} \beta_i < \alpha$, then it follows that $(\langle \alpha, x \rangle, [\beta^*, \gamma^*]) \in s$ for some γ^* .
- (5) If $\{(\langle \alpha_i, x_i \rangle, [\beta_i, \gamma_i]) : i < \omega\} \subset s$ and $\alpha^* = \sup_{i < \omega} \alpha_i$, then it follows that $(\langle \alpha^*, y \rangle, [\beta, \gamma]) \in s$ for some y, β, γ .
- (6) If $(\langle \alpha, x \rangle, [\beta, \bar{\gamma}]) \in s$ and $\text{cf}(\alpha) = \omega$, then $\{\beta < \alpha : \exists \gamma, (\langle \alpha, x \rangle, [\beta, \gamma]) \in s\}$ is unbounded in α .
- (7) If $(\langle \alpha, x \rangle, [\beta, \gamma]) \in s$ and $\text{cf}(\beta) = \omega$, then one of the following holds:
 - (a) there exists a sequence of ordinals $\langle \beta_i : i < \omega \rangle$ converging to β such that $\forall i < \omega, \exists \gamma_i, (\langle \alpha, x \rangle, [\beta_i, \gamma_i]) \in s$.
 - (b) there is some $\bar{\beta} < \beta$ and some sequence of ordinals $\langle \gamma_i : i < \omega \rangle$ converging to β such that $\forall i < \omega, (\langle \alpha, x \rangle, [\bar{\beta}, \gamma_i]) \in s$.
- (8) Suppose that $(\langle \alpha, x \rangle, [\beta, \gamma]) \in s$ and either $\text{cf}(\beta) = \omega_1$ or else $\{\beta' < \beta : \exists \gamma', (\langle \alpha, x \rangle, [\beta', \gamma']) \in s\}$ is unbounded in β . Then there is some $y \in \omega$ such that for all $\gamma' < \beta$, $(\langle \alpha, x \rangle, [\beta', \gamma']) \in s$ if and only if $(\langle \beta, y \rangle, [\beta', \gamma']) \in s$.

Suppose $s \in \mathbb{S}$ and there are α, β with $\beta < \alpha$ and $x, y \in \omega$ such that:

- (i) there are $\beta', \gamma', \beta'', \gamma''$ such that $(\langle \alpha, x \rangle, [\beta', \gamma']), (\langle \beta, y \rangle, [\beta'', \gamma'']) \in s$;
- (ii) for all $\gamma' < \beta$, $(\langle \alpha, x \rangle, [\beta', \gamma']) \in s$ if and only if $(\langle \beta, y \rangle, [\beta', \gamma']) \in s$.

Then we denote this $\text{Coh}(s, \alpha, x, \beta, y)$.

The ordering is defined so that $s' \leq_{\mathbb{S}} s$ holds if and only if:

- (1) $s' \supseteq s$.
- (2) If $\text{Coh}(s, \alpha, x, \beta, y)$ holds then $\text{Coh}(s', \alpha, x, \beta, y)$ holds.

We will also use the following conventions:

- An element of \mathfrak{S} is called a *bit*.
- If $s \in \mathbb{S}$, the set $\{\alpha < \aleph_2 : \exists x, \beta, \gamma, (\langle \alpha, x \rangle, [\beta, \gamma]) \in s\}$ is called the *domain* of s and is abbreviated $\text{dom}(s)$.
- If $s \in \mathbb{S}$, then $\text{dom}(s)$ has a maximal element γ , which will be denoted $\max(s)$.
- For $\langle \alpha, x \rangle \in \text{Lim}(\aleph_2) \times \omega$, we let

$$\dot{\mathcal{C}}_{\alpha}^x := \{(\check{\beta}, s) : \beta \geq 0, \exists \gamma, (\langle \alpha, x \rangle, [\beta, \gamma]) \in s \in \mathbb{S}\}.$$

³We are abusing the term “ordinal” in the case that $\beta = -1$.

- We sometimes use the terminology of coherence before we have verified that some $s \in \mathfrak{S}$ is in fact a condition in \mathbb{S} . The second point of the definition of $\leq_{\mathbb{S}}$ states that $s' \leq_{\mathbb{S}} s$ means that if s forces that y witnesses coherence of \dot{C}_α^x at β , then so does s' .
- If $\delta \in (\aleph_2 \cap \text{cof}(\omega_1)) \cup \{\aleph_2\}$ and $s \in \mathbb{S}$, we write $s \upharpoonright \delta := \{(\langle \alpha, x \rangle, [\beta, \gamma]) \in s : \alpha < \delta\}$, noting that $s \upharpoonright \delta \in \mathbb{S}$.
- For $s \in \mathbb{S}$, out $s = \{\beta \geq 0 : \exists \alpha, x, \gamma, (\langle \alpha, x \rangle, [\beta, \gamma]) \in s, s \Vdash \text{“ot}(\dot{C}_\alpha^x) = \omega\text{”}\}$.

We can describe an intuition for Definition 11 here. If $(\langle \alpha, x \rangle, [\beta, \gamma]) \in s \in \mathbb{S}$ and $\beta \geq 0$, then this indicates that s forces β to be a point in \dot{C}_α^x and also forces “ $\dot{C}_\alpha^x \cap (\beta, \gamma] = \emptyset$ ”. Note that x is just a placeholder label here. Clause 1 and Clause 2 should be thought of in association with Abraham’s presentation of Baumgartner’s forcing for adding a club in \aleph_1 with finite conditions [1], but generalized to \aleph_2 . Clause 3 asserts that if $\text{cf}(\alpha) = \omega_1$, then the set of \dot{C}_α^x for $x \in \omega$ is in fact a singleton. Clause 4 ensures that the clubs in the $\square(\aleph_2, \aleph_0)$ -sequence are closed. Clause 5 ensures that we will be able to extend any condition. Clause 6 ensures that the clubs in ordinals α of countable cofinality are unbounded. Clause 7 in part ensures that the question of whether β is a limit point of its club is determined by the condition, so that in closure arguments we only need to deal with freshly added limits of β ’s. Finally, Clause 8 ensures that the generic object added by \mathbb{S} is a coherent sequence. The second point of the definition of $\leq_{\mathbb{S}}$ will be used in the proof of Lemma 17 below.

The next steps are to establish that facts about \mathbb{S} that do not require countable closure.

Proposition 12. *Given $s \in \mathbb{S}$, $\alpha \in \text{dom } s$, and $x \in \omega$, the following are equivalent:*

- (1) $s \Vdash \text{“}\beta \in \lim \dot{C}_\alpha^x\text{”}$.
- (2) *Either:*
 - (a) $\text{cf}(\beta) = \omega_1$ and $(\langle \alpha, x \rangle, [\beta, \gamma]) \in s$ for some γ , or else
 - (b) there are sequences $\langle \beta_i : i < \omega \rangle$ and $\langle \gamma_i : i < \omega \rangle$ converging to β such that $(\langle \alpha, x \rangle, [\beta_i, \gamma_i]) \in s$ for all $i < \omega$.

Proof. (2) \implies (1): The implication is clear if (b) holds, so assume that (a) holds.

We consider the case $\text{cf}(\alpha) = \omega$. Suppose $s' \leq s$ (we are now suppressing the notation $\leq_{\mathbb{S}}$) and $\bar{\beta} < \beta$, where we want to show that there is some $s'' \leq s'$ such that $s'' \Vdash \text{“}[\bar{\beta}, \beta) \cap \dot{C}_\alpha^x \neq \emptyset\text{”}$.

Let

$$\beta^* = \sup\{\beta' < \beta : \exists \gamma', (\langle \alpha, x \rangle, [\beta', \gamma']) \in s'\},$$

so $\beta^* < \beta$ by the fact that s' is countable and $\text{cf}(\beta) = \omega_1$. Moreover, $\beta^* \in \text{dom}(s')$ by Clause 4.

If $\bar{\beta} \leq \beta^*$ then we are done. Otherwise, let

$$s'' = s' \cup \{(\langle \alpha, x \rangle, [\beta^*, \bar{\beta}])\} \cup \{(\langle \alpha, x \rangle, [\bar{\beta} + 1, \bar{\beta} + 1])\}.$$

Let us argue that s'' is a condition in the case that $\text{cf}(\alpha) = \omega$ by going through the clauses.

Clause 1: This is immediate since we are only adding two bits.

Clause 2: We do not need to consider bits of the form $(\langle \alpha', y \rangle, [\beta^*, \gamma^*])$ if $\alpha' \neq \alpha$. We also do not need to consider pairs of bits that are taken from s' . The clause clearly holds if both bits under consideration are the newly-added ones. The remaining cases take the form where one bit takes the form $(\langle \alpha, x \rangle, [\beta^*, \gamma^*]) \in s'$ and the other bit is one of the new ones, so the clause holds because we know that $\gamma^* < \beta^*$.

Clause 3: This holds vacuously since $\text{cf}(\alpha) = \omega$.

Clause 4: Any infinite sequence of bits under consideration will have an infinite subsequence in s' , so this clause holds because s' is a condition.

Clause 5: This holds for the same reason as Clause 4.

Clause 6: This holds because s' is a condition and

$$\{\langle \alpha, x \rangle : \exists \beta, \gamma, (\langle \alpha, x \rangle, [\beta, \gamma]) \in s'\} = \{\langle \alpha, x \rangle : \exists \beta, \gamma, (\langle \alpha, x \rangle, [\beta, \gamma]) \in s''\}.$$

Clause 7: The notable case is that in which we consider a bit of the form $(\langle \alpha, x \rangle, [\beta', \gamma']) \in s''$ and $\beta' = \bar{\beta} + 1$, in which case the clause holds vacuously. The other cases follow from the fact that s' is a condition.

Clause 8: If we are considering some $(\langle \alpha', x' \rangle, [\beta', \gamma']) \in s''$ and want to show that $\text{Coh}(s'', \alpha', x', \beta, y')$ holds for some $y' < \omega$, then this will hold because s'' is a condition, since the only relevant bits will be of the form $(\langle \alpha'', x'' \rangle, [\beta'', \gamma'']) \in s''$ where $\beta'' < \gamma'' < \beta^*$.

Given that s'' is a condition, it is clear that $s'' \Vdash "[\bar{\beta}, \beta) \cap \dot{C}_\alpha^x \neq \emptyset"$.

We also need to have $s'' \leq_{\mathbb{S}} s'$. We have $s'' \supseteq s'$, so we then need to consider the coherence clause for $\leq_{\mathbb{S}}$. This holds for reasons similar to the fact that Clause 8 holds in that no bits have been added where coherence needs to be verified.

The only substantive difference in the case where $\text{cf}(\alpha) = \omega_1$ is for Clause 3, which holds because it is explicitly taken care of for all $y \in \omega$. The argument that $s'' \leq_{\mathbb{S}} s'$ is also analogous in this case.

(1) \implies (2): Assume that β is a limit ordinal and that (2) does not hold. First suppose that $\text{cf}(\alpha) = \omega$. Let

$$\beta^* = \sup\{\beta' : \beta' < \beta : \exists \gamma', (\langle \alpha, x \rangle, [\beta', \gamma']) \in s'\}$$

if this set is not empty, otherwise $\beta^* = 0$. Then $\beta^* < \beta$, either by Clause 4 or by $\text{cf}(\beta) = \omega_1$.

If there is no γ such that $(\langle \alpha, x \rangle, [\beta, \gamma]) \in s$, then we let $s' = s \cup \{(\langle \alpha, x \rangle, [\beta^*, \beta])\}$. The argument that s' is a condition and that $s' \leq_{\mathbb{S}} s$ is similar to the backwards direction of the proof of this proposition, but it is strictly easier since we are only adding one bit. Then we see that $s' \Vdash "\dot{C}_\alpha^x \cap (\beta^*, \beta) = \emptyset"$ and hence $s' \Vdash "\beta \notin \lim \dot{C}_\alpha^x"$.

If there is some γ such that $(\langle \alpha, x \rangle, [\beta, \gamma]) \in s$, then $\text{cf}(\beta) = \omega$ (since we are assuming (2) does not hold). Then case (b) of Clause 7 holds, so there is some $\bar{\beta} < \beta$ and a sequence of γ_i 's such that $(\langle \alpha, x \rangle, [\bar{\beta}, \gamma_i]) \in s$ for all $i < \omega$. Hence $s \Vdash "\dot{C}_\alpha^x \cap (\bar{\beta}, \beta) = \emptyset"$ and therefore $s \Vdash "\beta \notin \lim \dot{C}_\alpha^x"$.

An analogous argument applies for the case that $\text{cf}(\alpha) = \omega_1$. \square

Proposition 13. *The name \dot{C}_α^x is forced to be a closed unbounded set in α for all limit ordinals $\alpha < \aleph_2$ and all $x < \omega$.*

Proof. First we show that the \dot{C}_α^x 's are forced to be closed. (We will prove nonemptiness when we prove unboundedness.) If there are β' and γ' with $(\langle \alpha, x \rangle, [\beta', \gamma']) \in \bar{s}$, then \bar{s} forces \dot{C}_α^x to be closed under countable sequences by Clause 4. Closure under uncountable sequences follows from Proposition 12 since if $s \in \mathbb{S}$ forces $\beta \notin \lim \dot{C}_\alpha^x$ for some β with $\text{cf}(\beta) = \omega_1$, then s forces $\beta \notin \dot{C}_\alpha^x$.

Most of the proof then consists of showing that the \dot{C}_α^x 's are forced to be unbounded and in particular nonempty.

Consider the case of α with $\text{cf}(\alpha) = \omega$. Fix some $\bar{s} \in \mathbb{S}$. If there are β and γ with $(\langle \alpha, x \rangle, [\beta, \gamma]) \in \bar{s}$, then \bar{s} forces \dot{C}_α^x to be unbounded by Clause 6.

Now suppose there are no β and γ with $(\langle \alpha, x \rangle, [\beta, \gamma]) \in \bar{s}$. Take a sequence $\langle \delta_i : i < \omega \rangle$ converging to α where $\delta_0 = 0$ and δ_i is a successor ordinal for $i > 0$. Then let $s := \bar{s} \cup \{(\langle \alpha, x \rangle, [\delta_i, \delta_{i+1} - 1]) : x \in X, i < \omega\}$. As long as we show that s is a condition such that $s \leq_{\mathbb{S}} \bar{s}$, it is clear that s forces \dot{C}_α^x to be unbounded and (vacuously) closed in α .

We argue that s is a condition, going through the less trivial clauses:

Clause 2: The intervals $[\delta_i, \delta_{i+1} - 1]$ are pairwise disjoint.

Clause 4: This closure holds vacuously when the new bits are considered.

Clause 6: The δ_i 's are unbounded in α .

Clause 8: The only notable case is where $\bar{s} \Vdash \text{“}\alpha \in \lim \dot{C}_{\alpha'}^{x'}\text{”}$ for some $\langle \alpha', x' \rangle$, but in this case we would already have $\text{Coh}(\bar{s}, \alpha', x', \beta, y)$ for some $y < \omega$.

The fact that $s' \leq_{\mathbb{S}} s$ is quite immediate given the argument for Clause 8.

Now we consider the case of α with $\text{cf}(\alpha) = \omega_1$.

For the case where $\text{cf}(\alpha) = \omega_1$. Fix $x < \omega$ and $\bar{\beta} < \alpha$ and some $\bar{s} \in \mathbb{S}$ such that for some β', γ' , $(\langle \alpha, x \rangle, [\beta', \gamma']) \in \bar{s}$.

Let

$$\beta^* = \sup\{\beta' < \alpha : \exists \gamma', (\langle \alpha, x \rangle, [\beta', \gamma']) \in \bar{s}\},$$

so $\beta^* < \alpha$ by the fact that s' is countable and $\text{cf}(\beta) = \omega_1$. If $\beta^* \geq \bar{\beta}$ then we are done. (We can also allow the set to be empty and for β^* to be 0.)

Now let

$$s = \bar{s} \cup \{(\langle \alpha, y \rangle, [\beta^*, \bar{\beta}]) : y < \omega\} \cup \{(\langle \alpha, y \rangle, [\bar{\beta} + 1, \bar{\beta} + 1]) : y < \omega\}.$$

The proof that s is a condition such that $s \leq_{\mathbb{S}} \bar{s}$ is very similar to the first part of the proof of Proposition 12. From there it is immediate that $s \Vdash \text{“}\dot{C}_\alpha^x \cap (\bar{\beta}, \alpha) \neq \emptyset\text{”}$. \square

Now we will work towards proving that a dense subset of \mathbb{S} is countably closed. First we define the subset:

Definition 14. We say that $s \in \mathbb{S}$ is *complete* if the following holds: If $(\langle \alpha, x \rangle, [\beta, \gamma]) \in s$ and there are $\alpha' < \alpha$ and some $x' \in \omega$ such that $(\langle \alpha', x' \rangle, [\beta', \gamma']) \in s$, then:

- there are $\beta'' \leq \alpha'$ and $\gamma'' \geq \alpha'$ such that $(\langle \alpha, x \rangle, [\beta'', \gamma'']) \in s$,
- there are $\beta''' \leq \beta'$ and $\gamma''' \geq \beta'$ such that $(\langle \alpha, x \rangle, [\beta''', \gamma''']) \in s$.

Proposition 15. *The set of complete conditions is dense in \mathbb{S} . More precisely, if $\bar{s} \in \mathbb{S}$, then there is a complete $s \leq_{\mathbb{S}} \bar{s}$ such that $\text{dom } s = \text{dom } \bar{s}$.*

Proof. Fix $\bar{s} \in \mathbb{S}$. For all $\alpha \in \text{dom } \bar{s}$ and $x < \omega$, define

$$B = \{\beta : \exists \alpha, x, \gamma, (\langle \alpha, x \rangle, [\beta, \gamma]) \in \bar{s}\}$$

and

$$B_{\langle \alpha, x \rangle} = \{\beta : \exists \gamma, (\langle \alpha, x \rangle, [\beta, \gamma]) \in \bar{s}\}$$

and let

$$s = \bar{s} \cup \{(\langle \alpha, x \rangle, [\beta, \gamma]) : \exists \beta', \gamma', (\langle \alpha, x \rangle, [\beta', \gamma']) \in \bar{s}, \\ \gamma \in \text{dom } \bar{s} \cup B, \beta = \sup(B_{\langle \alpha, x \rangle} \cap \gamma)\}.$$

We can see that s is complete by construction. Now we verify that s is a condition.

Clause 1: Immediate.

Clause 2: Fix α and x for consideration. Suppose $(\langle \alpha, x \rangle, [\beta, \gamma]) \in s$ and $(\langle \alpha, x \rangle, [\beta', \gamma']) \in s$. If they are both in \bar{s} then we are done. If they are both in $s \setminus \bar{s}$ with $\beta \leq \beta'$, then by definition there are no points in $B_{\langle \alpha, x \rangle}$ between β and γ , so either $\beta = \beta'$ or $\gamma < \beta'$. If without loss of generality the first of these bits is in \bar{s} and the other is in $s \setminus \bar{s}$ with $\beta \leq \beta'$, then we know that there are no elements of $B_{\langle \alpha, x \rangle}$ in the interval $(\beta, \gamma]$, so either $\beta = \beta'$ or $\gamma < \beta'$. If we assume that $\beta' \leq \beta$ the argument is similar.

Clause 3: This property is inherited by s from \bar{s} because the same changes are made to every \dot{C}_α^x for each $x < \omega$.

Clause 4: We have for all α and x that

$$\{\beta : \exists \gamma, (\langle \alpha, x \rangle, [\beta, \gamma]) \in s\} = \{\beta : \exists \gamma, (\langle \alpha, x \rangle, [\beta, \gamma]) \in \bar{s}\}$$

and therefore the fact that Clause 4 holds for s follows from the fact that it holds for \bar{s} .

Clause 5: This holds because $\text{dom } \bar{s} = \text{dom } s$ and moreover the x 's for which \dot{C}_α^x is defined are the same.

Clause 6: Holds for the same reason as Clause 5.

Clause 7: Holds for the same reason as Clause 4.

Clause 8: Fix $\langle \alpha, x \rangle$. Suppose first that $\{\beta' : \exists \gamma', (\langle \alpha, x \rangle, [\beta', \gamma']) \in s\}$ is unbounded in β . Then it follows that $\{\beta' : \exists \gamma', (\langle \alpha, x \rangle, [\beta', \gamma']) \in \bar{s}\}$ is unbounded in β and therefore $\text{Coh}(\bar{s}, \alpha, x, \beta, y)$ holds for some $y < \omega$. But if we write

$$B' = \{\beta' < \beta : \exists \alpha, x, \gamma', (\langle \alpha, y \rangle, [\beta', \gamma']) \in \bar{s}\}$$

then $B \cap \beta = B'$. Moreover, $B_{\langle \alpha, x \rangle} \cap \beta = B_{\langle \beta, y \rangle}$. Therefore, given how the bits in \bar{s} are defined, we have $\text{Coh}(s, \alpha, x, \beta, y)$.

Finally, see that $s \leq_{\mathbb{S}} \bar{s}$, which follows because clearly $s \supseteq \bar{s}$ and by an argument analogous to the argument for Clause 8. \square

The proof that the set of complete conditions in \mathbb{S} is countably closed is similar to the analogous proof for Jensen's method of forcing square sequences in the sense that ω -sequences can vacuously top off lower bounds of countable sequences. However, it is different in the sense that new limit

points of the \dot{C}_α^x 's are added. Because of this complication, we introduce some additional terminology.

Definition 16. Let $\vec{s} = \langle s_i : i < \omega \rangle$ be an $\leq_{\mathbb{S}}$ -decreasing sequence of conditions.

- Let $\text{ed}(\vec{s}) = \bigcup_{i < \omega} \text{dom}(s_i)$ be the *existing domain*.
- Let $\text{fd}(\vec{s}) = \lim^+(\text{ed}(\vec{s})) \setminus \text{ed}(\vec{s})$ be the *fresh domain*.
- Let the *existing points* $\text{ep}(\vec{s})$ consist of $\beta \in \aleph_2$ such that for some $\alpha \in \text{ed}$, $x \in \omega$, and $\gamma < \aleph_2$, $(\langle \alpha, x \rangle, [\beta, \gamma]) \in \bigcup_{i < \omega} s_i$.
- Suppose $\beta \in \lim(\aleph_2)$ and $\alpha \in \text{ed}(\vec{s})$ as witnessed by $(\langle \alpha, x \rangle, [\beta', \gamma'])$. Suppose that $\beta \notin \text{ep}(\vec{s})$ and $\beta < \alpha$, but there are sequences $\langle \beta_n : n < \omega \rangle$ and $\langle \gamma_n : n < \omega \rangle$ with supremum β such that $\forall n < \omega$, $(\langle \alpha, x \rangle, [\beta_n, \gamma_n]) \in \bigcup_{i < \omega} s_i$. Then we let $\beta \in \text{fp}(\vec{s})$, the set of *fresh points*, and we say that $\langle \alpha, x \rangle$ *witnesses* $\beta \in \text{fp}(\vec{s})$.

If \bar{s} is a lower bound of \vec{s} , then we say that \bar{s} is *parsimonious* if:

- $\text{dom}(\bar{s}) = \text{ed}(\vec{s}) \cup \text{fd}(\vec{s}) \cup \text{fp}(\vec{s})$;
- for all $\alpha \in \text{Lim}(\aleph_2)$ and $x \in \omega$, $(\langle \alpha, x \rangle, [\beta, \gamma]) \in \bar{s}$ implies either $\beta \in \text{ep}(\vec{s}) \cup \text{fp}(\vec{s})$ or $\bar{s} \Vdash \text{“ot}(\dot{C}_\alpha^x) = \omega\text{”}$.

We drop the notation for \vec{s} when the context is clear.

Lemma 17. *The set of complete conditions in \mathbb{S} is countably closed. Moreover, every $\leq_{\mathbb{S}}$ -decreasing sequence $\langle s_i : i < \omega \rangle$ of complete conditions has a parsimonious lower bound.*

Before proving the lemma, we establish the basic relationships between the sets of points described above.

Proposition 18. *Let $\vec{s} = \langle s_i : i < \omega \rangle$ be an $\leq_{\mathbb{S}}$ -decreasing sequence of complete conditions in \mathbb{S} .*

- (1) $\text{fd}(\vec{s}), \text{fp}(\vec{s}) \subset \aleph_2 \cap \text{cof}(\omega)$.
- (2) $\text{fp}(\vec{s}) \cap \text{ep}(\vec{s}) = \text{fp}(\vec{s}) \cap \text{ed}(\vec{s}) = \emptyset$.

Proof. (1) is immediate, so we prove (2). Suppose firstly that $\beta \in \text{fp}(\vec{s}) \cap \text{ep}(\vec{s})$ and that in particular $\beta \in \text{fp}(\vec{s})$ is witnessed by $\langle \alpha, x \rangle$. Since $\beta \in \text{ed}(\vec{s})$ there is some $i < \omega$ and some α, x, γ such that $(\langle \alpha, x \rangle, [\beta, \gamma]) \in \text{dom } s_i$. But by completeness of s_i and the fact that $\beta < \alpha$, we have some β' and γ' such that $(\langle \alpha, x \rangle, [\beta', \gamma']) \in s_i$ and $\beta' \leq \beta \leq \gamma'$. But then this contradicts $\beta \in \text{fp}(\vec{s})$. The argument that $\text{fp}(\vec{s}) \cap \text{ed}(\vec{s}) = \emptyset$ is analogous. \square

Proof of Lemma 17. Let $\langle s_i : i < \omega \rangle$ be an $\leq_{\mathbb{S}}$ -decreasing sequence of complete conditions in \mathbb{S} and fix ed , fd , ep , and fp as in Definition 16. Let $S_0 := \bigcup_{i < \omega} s_i$. For each $\beta \in \text{fp}$, let X_β be the set of pairs $\langle \alpha, x \rangle$ witnessing that $\beta \in \text{fp}$. We let

$$S_1 := \{(\langle \alpha, x \rangle, [\beta, \beta]) : \alpha \in \text{ed}, \beta \in \text{fp}, \langle \alpha, x \rangle \in X_\beta\}.$$

For each $\beta \in \text{fp}$, let $f_\beta : \omega \rightarrow X_\beta$ be a surjection. Let

$$S_2 := \{(\langle \beta, x \rangle, [\beta', \gamma']) : \beta \in \text{fp}, f_\beta(x) = \langle \alpha, y \rangle \in X_\beta, \\ (\langle \alpha, y \rangle, [\beta', \gamma']) \in S_0, \gamma' < \beta\}.$$

Also let

$$S_3 := \{(\langle \beta, x \rangle, [\beta', \beta']) : \beta' < \beta, \mathbf{fp} \supseteq \{\beta, \beta'\}, f_\beta(x) = \langle \alpha, y \rangle \in X_\beta \cap X_{\beta'}\}^4$$

For each $\alpha \in \mathbf{fd} \setminus \mathbf{fp}$, choose a sequence $\langle \delta_n^\alpha : n < \omega \rangle$ of ordinals converging to α where $\delta_0^\alpha = 0$ and δ_n^α is a successor ordinal for all $n > 0$. Let

$$S_4 := \{(\langle \alpha, x \rangle, [\delta_n^\alpha, \delta_{n+1}^\alpha - 1]) : \alpha \in \mathbf{fd} \setminus \mathbf{fp}, x \in \omega, n < \omega\}.$$

The lower bound we seek is $s := S_0 \cup S_1 \cup S_2 \cup S_3 \cup S_4$. It is clear that s is parsimonious as long as it is a lower bound. It remains to argue that s is a condition and that $s \leq_S s_i$ for all $i < \omega$.

Assuming that s is in fact a condition, it is relatively straightforward to argue that it is a lower bound: Since we have $s \supseteq s_i$ for all $i < \omega$, the first point in the definition of \leq_S holds. For the second point, suppose that $\text{Coh}(s_i, \alpha, x, \beta, y)$ holds. Then $\alpha, \beta \in \mathbf{ed}$. Since Clause 8 holds for s_i we do not need to worry about bits from S_0 , for which Clause 8 is witnessed by other bits from S_0 , and so we only need to consider bits from S_1 since this is the only one of the S_k 's with bits of the form $(\langle \alpha^*, x^* \rangle, [\beta^*, \gamma^*])$ with $\alpha^* \in \mathbf{ed}$. Suppose that $(\langle \alpha, x \rangle, [\beta', \beta']) \in S_1$ with $\beta' < \beta$, i.e. $\beta' \in \mathbf{fp} \cap \beta$, and $\langle \alpha, x \rangle \in X_{\beta'}$. Then $\langle \beta, y \rangle \in X_{\beta'}$ as well, so $(\langle \beta, y \rangle, [\beta', \beta']) \in S_1$. The reverse reasoning in which we start by considering $(\langle \beta, y \rangle, [\beta', \beta']) \in S_1$ also holds.

For proving that s is a condition, we first observe that if $(\langle \alpha, x \rangle, [\beta, \gamma]) \in s$, then exactly one of the following holds: $\alpha \in \mathbf{ep}$, $\alpha \in \mathbf{fd} \setminus \mathbf{fp}$, and $\alpha \in \mathbf{fp}$. This is because it follows from the definitions that either $\alpha \in \mathbf{ep}$ or $\alpha \in \mathbf{fp}$, and we have that $\alpha \in \mathbf{ep} \cap \mathbf{fp} = \emptyset$ by Proposition 18. Moreover, exactly one of the following holds: $\beta \in \mathbf{ep}$ or $\beta \in \mathbf{fp}$, also by Proposition 18.

Clause 1: This follows because the S_k 's are each defined from countably many parameters.

Clause 2: Fix $\langle \alpha, x \rangle$, $[\beta, \gamma]$, and $[\beta', \gamma']$. We consider the possible cases:

$\alpha \in \mathbf{ed}$, $\beta, \beta' \in \mathbf{ep}$: If i is large enough that $(\langle \alpha, x \rangle, [\beta, \gamma])$ and $(\langle \alpha, x \rangle, [\beta', \gamma'])$ are in s_i , then this follows from the fact s_i is a condition.

$\alpha \in \mathbf{ed}$, $\beta \in \mathbf{ep}$, $\beta' \in \mathbf{fp}$: Then $\gamma' = \beta'$, so we can assume $\beta < \beta'$ and we want to show that $\gamma < \beta$. If this were not the case, then because of the fact that we have some $i < \omega$ with $(\langle \alpha, x \rangle, [\beta, \gamma]) \in s_i$, and because $\langle \alpha, x \rangle$ would witness $\beta' \in \mathbf{fp}$ in this case, it follows that $\beta \leq \gamma$ would imply that $\beta' \notin \mathbf{fp}$, a contradiction.

$\alpha \in \mathbf{ed}$, $\beta', \beta \in \mathbf{fp}$: Then $\gamma' = \beta'$ and $\gamma = \beta$, so there is nothing to deal with.

$\alpha \in \mathbf{fd} \setminus \mathbf{fp}$: Then the clause follows from the fact that the intervals $[\delta_n^\alpha, \delta_{n+1}^\alpha - 1]$ are disjoint.

$\alpha \in \mathbf{fp}$, $\beta, \beta' \in \mathbf{ep}$: Then there is some $\tilde{x} \in \omega$ and $\tilde{\alpha} \in \mathbf{ep}$ such that $f_\alpha(x) = \langle \tilde{\alpha}, \tilde{x} \rangle$. Then $(\langle \alpha, x \rangle, [\beta, \gamma]), (\langle \alpha, x \rangle, [\beta', \gamma']) \in S_2$, and so the fact that we have $(\langle \tilde{\alpha}, \tilde{x} \rangle, [\beta, \gamma]), (\langle \tilde{\alpha}, \tilde{x} \rangle, [\beta', \gamma']) \in s_i$ for large enough i gives us the clause for this case.

$\alpha \in \mathbf{fp}$, $\beta \in \mathbf{ep}$, $\beta' \in \mathbf{fp}$: We can assume $\beta < \beta'$. It must be the case that $(\langle \alpha, x \rangle, [\beta', \beta']) \in S_3$ as witnessed by some $\langle \tilde{\alpha}, \tilde{x} \rangle$ with $f_\alpha(x) = \langle \tilde{\alpha}, \tilde{x} \rangle$ and moreover that $\langle \tilde{\alpha}, \tilde{x} \rangle \in X_{\beta'}$. Then $(\langle \alpha, x \rangle, [\beta, \gamma]) \in S_2$, so there is some

⁴This means that if $\langle \alpha, x \rangle$ witnesses that both $\beta \in \mathbf{fp}$ and $\beta' \in \mathbf{fp}$ where $\beta' < \beta$, we make sure to put the point β' in \dot{C}_β^α where we will show that $\text{Coh}(s, \alpha, y, \beta, x)$ holds.

$i < \omega$ such that $(\langle \tilde{\alpha}, \tilde{x} \rangle, [\beta, \gamma]) \in s_i$. Then if it were the case that $\beta' \leq \gamma$, it would not be the case that $\langle \tilde{\alpha}, \tilde{x} \rangle \in X_{\beta'}$.

$\alpha \in \mathbf{fp}$, $\beta, \beta' \in \mathbf{fp}$: This is analogous to the case $\alpha \in \mathbf{ed}$, $\beta, \beta' \in \mathbf{fp}$ in the sense that we only need to consider whether or not $\beta = \beta'$.

Clause 3: If $\text{cf}(\alpha) = \omega_1$ and $\alpha \in \text{dom}(\bar{s})$, then $\alpha \in \mathbf{ed}$. Then \bar{s} inherits Clause 3 from the fact that the s_i 's have Clause 3.

Clause 4: If $\alpha \in \mathbf{fd} \setminus \mathbf{fp}$ this clause holds vacuously. For the other cases, suppose that $\{(\langle \alpha, x \rangle, [\beta_n, \gamma_n]) : i < \omega\} \subseteq s$ and $\beta^* = \sup_{n < \omega} \beta_n < \alpha$. We will subdivide the cases based on the minimal value k for which infinitely many of these bits are taken from S_k .

Inf. many from S_0 : Choose $i < \omega$ large enough so that $(\langle \alpha, x \rangle, [\beta_n, \gamma_n]) \in s_i$ for some n in our infinite set. Then $(\langle \alpha, x \rangle, [\beta_n, \gamma_n]) \in s_i$ will hold for infinitely many n . Therefore we can see that either $\beta^* \in \mathbf{ep}$ or $\beta^* \in \mathbf{fp}$. If $\beta^* \in \mathbf{ep}$ then we are done. If $\beta^* \in \mathbf{fp}$, then this is witnessed by $\langle \alpha, x \rangle$, so we have $(\langle \alpha, x \rangle, [\beta^*, \beta^*]) \in S_1 \subseteq s$.

Inf. many from S_1 : Then we have infinitely many $(\langle \alpha, x \rangle, [\beta_n, \beta_n])$ where $\langle \alpha, x \rangle \in X_{\beta_n}$. So for each such $n < \omega$ there is a sequence β_n^j, γ_n^j converging to β_n such that $(\langle \alpha, x \rangle, [\beta_n^j, \gamma_n^j]) \in S_0$, so it follows that $\beta^* \in \mathbf{fp}$ and that $\langle \alpha, x \rangle \in X_{\beta^*}$. Therefore $(\langle \alpha, x \rangle, [\beta^*, \beta^*]) \in S_1 \subseteq s$.

Inf. many from S_2 : Then let $\langle \tilde{\alpha}, \tilde{x} \rangle$ be such that $f_\alpha(x) = \langle \tilde{\alpha}, \tilde{x} \rangle$. Then we are saying that we have infinitely many $n < \omega$ such that $(\langle \tilde{\alpha}, \tilde{x} \rangle, [\beta_n, \gamma_n]) \in S_0$. Therefore $\beta^* \in \mathbf{fp}$, as witnessed by $\langle \tilde{\alpha}, \tilde{x} \rangle$, and hence $(\langle \alpha, x \rangle, [\beta^*, \beta^*]) \in S_3 \subseteq s$.

Inf. many from S_3 : Again choose $\langle \tilde{\alpha}, \tilde{x} \rangle$ such that $f_\alpha(x) = \langle \tilde{\alpha}, \tilde{x} \rangle$. We are saying that we have infinitely many $n < \omega$ such that $(\langle \tilde{\alpha}, \tilde{x} \rangle, [\beta_n, \beta_n]) \in S_3$ where $\langle \tilde{\alpha}, \tilde{x} \rangle$ also witnesses that each β_n is in \mathbf{fp} . This implies that $\langle \tilde{\alpha}, \tilde{x} \rangle$ witnesses that β^* is in \mathbf{fp} . Hence $(\langle \alpha, x \rangle, [\beta^*, \beta^*]) \in S_3 \subseteq s$.

Inf. many from S_4 : We already ruled this case out.

Clause 5: Suppose $\{(\langle \alpha_i, x_i \rangle, [\beta_i, \gamma_i]) : i < \omega\} \subset s$ and $\alpha^* = \sup_{i < \omega} \alpha_i$. All possibilities are among the following:

$\alpha^* \in \mathbf{ed}$: Then there is some $i < \omega$ such that $\alpha^* \in \text{dom } s_i$, hence $\alpha^* \in \text{dom } s$.

$\alpha^* \in \mathbf{fd} \setminus \mathbf{fp}$: Then for all $n < \omega$ we have $(\langle \alpha^*, x \rangle, [\delta_n^\alpha, \delta_{n+1}^\alpha - 1]) \in S_4 \subseteq s$ and so we have $\alpha^* \in \text{dom } s$.

$\alpha^* \in \mathbf{fp}$: Then there is some $\langle \alpha, y \rangle$ witnessing $\alpha^* \in \mathbf{fp}$ and some x such that $f_{\alpha^*}(x) = \langle \alpha, y \rangle$. Choose some $(\langle \alpha, y \rangle, [\beta, \gamma]) \in S_0$ with $\gamma < \alpha^*$. Then $(\langle \alpha^*, x \rangle, [\beta, \gamma]) \in S_2 \subseteq s$.

Clause 6: This is clear by inspection.

Clause 7: If $\alpha \in \mathbf{fd} \setminus \mathbf{fp}$ then the clause holds vacuously.

For the other cases, suppose that $(\langle \alpha, x \rangle, [\beta, \gamma]) \in s$ and $\text{cf}(\beta) = \omega$.

$\alpha \in \mathbf{ed}, \beta \in \mathbf{ep}$: Then the clause holds because $(\langle \alpha, x \rangle, [\beta, \gamma]) \in s_i$ for large i and s_i is a condition.

$\alpha \in \mathbf{ed}, \beta \in \mathbf{fp}$: Then we have $(\langle \alpha, x \rangle, [\beta, \gamma]) \in S_1$ where $\beta = \gamma$ and $\langle \alpha, x \rangle$ witnesses $\beta \in \mathbf{fp}$. Then it follows that case (a) holds.

$\alpha \in \mathbf{fp}, \beta \in \mathbf{ep}$: In this case we have $(\langle \alpha, x \rangle, [\beta, \gamma]) \in S_2$, meaning that $f_\alpha(x) = \langle \tilde{\alpha}, \tilde{x} \rangle$ for some $\langle \tilde{\alpha}, \tilde{x} \rangle \in X_\alpha$, and that $(\langle \tilde{\alpha}, \tilde{x} \rangle, [\beta, \gamma]) \in S_0$. Hence if

$i < \omega$ is large enough that $(\langle \tilde{\alpha}, \tilde{x} \rangle, [\beta, \gamma]) \in s_i$, then the clause is witnessed by the fact that s_i is a condition and $\text{Coh}(s_i, \tilde{\alpha}, \tilde{x}, \alpha, x)$ holds.

$\alpha \in \mathbf{fp}, \beta \in \mathbf{fp}$: In this case we have $(\langle \alpha, x \rangle, [\beta, \gamma]) \in S_2$, meaning that $\beta = \gamma$, $f_\alpha(x) = \langle \tilde{\alpha}, \tilde{x} \rangle$, and $\langle \tilde{\alpha}, \tilde{x} \rangle$ also witnesses $\beta \in \mathbf{fp}$. Hence there are β_i, γ_i converging to β such that $(\langle \tilde{\alpha}, \tilde{x} \rangle, [\beta_i, \gamma_i]) \in S_0$ for $i < \omega$. Therefore $(\langle \alpha, x \rangle, [\beta_i, \gamma_i]) \in S_2$ for $i < \omega$. This implies that case (a) holds.

Clause 8: Referring to the statement in Definition 11, fix $\langle \alpha, x \rangle$ and β such that the hypothesis holds for $(\langle \alpha, x \rangle, [\beta, \gamma]) \in s$ for some γ .

$\alpha \in \mathbf{fd} \setminus \mathbf{fp}$: Then coherence holds vacuously because there are no limit points to consider.

$\alpha \in \mathbf{ed}, \beta \in \mathbf{ep}$: Since $\beta \in \mathbf{ep}$, there is some $i < \omega$ such that $s_i \Vdash \text{“}\beta \in \lim \dot{C}_\alpha^x\text{”}$. Therefore there is some y witnessing that $\text{Coh}(s_i, \alpha, x, \beta, y)$ holds. This also holds for $i' \geq i$ by the definition of \leq_s . This means that if $(\langle \alpha, x \rangle, [\beta', \gamma']) \in s$ for $\gamma' < \beta$, then there are two possibilities. The first is that $\beta' \in \mathbf{ep}$, and hence $(\langle \alpha, x \rangle, [\beta', \gamma']) \in s_j$ for some $j \geq i$, in which $\text{Coh}(s_j, \alpha, x, \beta, y)$ implies that $(\langle \beta, y \rangle, [\beta', \gamma']) \in s_j \subseteq s$. The second possibility is that $\beta' \in \mathbf{fp}$. Then it must be the case that $(\langle \alpha, x \rangle, [\beta', \gamma']) \in S_1$, meaning that $\beta' = \gamma'$ and that $\langle \alpha, x \rangle \in X_{\beta'}$. Then this plus the fact that $\text{Coh}(s_i, \alpha, x, \beta, y)$ holds for large i implies that $\langle \beta, y \rangle \in X_{\beta'}$. Therefore $(\langle \beta, y \rangle, [\beta', \beta']) \in S_1$.

Now suppose $(\langle \beta, y \rangle, [\beta', \gamma']) \in s$ and consider the same two possibilities. If $\beta' \in \mathbf{ep}$, then if i is large enough that $s_i \Vdash \text{“}\beta \in \lim \dot{C}_\alpha^x\text{”}$ and $(\langle \beta, y \rangle, [\beta', \gamma']) \in s_i$, then $\text{Coh}(s_i, \alpha, x, \beta, y)$ implies $(\langle \alpha, x \rangle, [\beta', \gamma']) \in s_i$. If $\beta' \in \mathbf{fp}$ then $(\langle \beta, y \rangle, [\beta', \gamma']) \in S_1$, implying $\beta' = \gamma'$ and $\langle \beta, y \rangle \in X_{\beta'}$, so $\langle \alpha, x \rangle \in X_{\beta'}$, hence $(\langle \alpha, x \rangle, [\beta', \gamma']) \in S_1$.

$\alpha \in \mathbf{ed}, \beta \in \mathbf{fp}$: Then $\langle \alpha, x \rangle$ witnesses $\beta \in \mathbf{fp}$, so let y be such that $f_\beta(y) = \langle \alpha, x \rangle$. Then we can argue that $\text{Coh}(s, \alpha, x, \beta, y)$ holds. If $(\langle \alpha, x \rangle, [\beta', \gamma']) \in s$ and $\beta' \in \mathbf{ep}$, then $(\langle \alpha, x \rangle, [\beta', \gamma']) \in S_0$, and hence $(\langle \beta, y \rangle, [\beta', \gamma']) \in S_2$. If $(\langle \alpha, x \rangle, [\beta', \gamma']) \in s$ and $\beta' \in \mathbf{fp}$, then $\gamma' = \beta'$ and $\langle \alpha, x \rangle \in X_{\beta'}$, so $(\langle \beta, y \rangle, [\beta', \gamma']) \in S_3$. The other inclusion follows the same reasoning.

$\alpha \in \mathbf{fp}, \beta \in \mathbf{ep}$: Then there is some $\langle \tilde{\alpha}, \tilde{x} \rangle$ witnessing $\alpha \in \mathbf{fp}$ with $f_\alpha(x) = \langle \tilde{\alpha}, \tilde{x} \rangle$. All bits $(\langle \alpha, x \rangle, [\beta, \gamma]) \in s$ such that $\beta \in \mathbf{ep}$ are in S_2 and witnessed by $(\langle \tilde{\alpha}, \tilde{x} \rangle, [\beta, \gamma]) \in S_0$. Therefore there is some $i < \omega$ with $s_i \Vdash \text{“}\beta \in \lim \dot{C}_\alpha^{\tilde{x}}\text{”}$. Let y be such that $\text{Coh}(s_i, \tilde{\alpha}, \tilde{x}, \beta, y)$ holds. We can then argue that $\text{Coh}(s, \alpha, x, \beta, y)$ holds. Suppose that $\beta', \gamma' < \beta$ and $(\langle \alpha, x \rangle, [\beta', \gamma']) \in \tilde{S}$ with $\beta' \in \mathbf{ep}$. Then $(\langle \tilde{\alpha}, \tilde{x} \rangle, [\beta', \gamma']) \in S_0$, so $(\langle \beta, y \rangle, [\beta', \gamma']) \in S_0$. If instead $\beta' \in \mathbf{fp}$, then $(\langle \alpha, x \rangle, [\beta', \beta']) \in S_3$ and $\langle \tilde{\alpha}, \tilde{x} \rangle \in X_{\beta'}$, which implies $\langle \beta, y \rangle \in X_{\beta'}$, so $(\langle \beta, y \rangle, [\beta', \beta']) \in S_3$. Again, the reverse reasoning applies if we start with the premise that $(\langle \beta, y \rangle, [\beta', \gamma']) \in s$.

$\alpha \in \mathbf{fp}, \beta \in \mathbf{fp}$: Then $(\langle \alpha, x \rangle, [\beta, \beta]) \in S_3$, so there is some $\langle \tilde{\alpha}, \tilde{x} \rangle \in X_\alpha \cap X_\beta$ such that $f_\alpha(x) = \langle \tilde{\alpha}, \tilde{x} \rangle$. Let y be such that $f_\beta(y) = \langle \tilde{\alpha}, \tilde{x} \rangle$. We argue that $\text{Coh}(s, \alpha, x, \beta, y)$ holds. Suppose $(\langle \alpha, x \rangle, [\beta', \gamma']) \in s$. If $\beta' \in \mathbf{ep}$, then $(\langle \alpha, x \rangle, [\beta', \gamma']) \in S_2$, so $(\langle \tilde{\alpha}, \tilde{x} \rangle, [\beta', \gamma']) \in S_0$, so $(\langle \beta, y \rangle, [\beta, \gamma']) \in S_2$. If $\beta' \in \mathbf{fp}$, then $\beta' = \gamma'$ and $(\langle \alpha, x \rangle, [\beta', \beta']) \in S_3$, implying $\langle \tilde{\alpha}, \tilde{x} \rangle \in X_{\beta'} \cap X_\alpha$, i.e. $\langle \tilde{\alpha}, \tilde{x} \rangle \in X_{\beta'} \cap X_\beta \cap X_\alpha$, implying $(\langle \beta, y \rangle, [\beta', \beta']) \in S_3$ also. Once more, we can reverse the reasoning.

This finishes the proof of Lemma 17. \square

The next task is to prove that \mathbb{S} preserves \aleph_2 .

Lemma 19. *If $s \in \mathbb{S}$ and $M \prec K$ is a basic model with $s, \mathbb{S} \in M$, then s is strongly (M, \mathbb{S}) -generic.*

Therefore Proposition 6 implies:

Corollary 20. *Under CH, \mathbb{S} preserves \aleph_2 .*

Proof of Lemma 19. Let $\delta = M \cap \aleph_2$. Suppose $s' \leq s$ and let s^* consist of all $(\langle \alpha, x \rangle, [\beta, \gamma]) \in s'$ such that $\alpha, \beta, \gamma < \delta$. Then $s^* \in M$ by the countable closure of M . Let $s^{**} \leq s^*$ where $s^{**} \in M$. We will construct s'' such that $s'' \leq s', s^{**}$.

For each $\alpha \in \text{dom}(s') \setminus \delta$, $x \in \omega$, let $X_{\langle \alpha, x \rangle}$ be the set of $\langle \beta, y \rangle$ such that $s' \Vdash \text{“}\beta \in \lim \dot{C}_\alpha^x \cap \delta\text{”}$, and such that y witnesses coherence, i.e. Clause 8. Let S_0 be the set

$$\{(\langle \alpha, x \rangle, [\beta', \gamma']) : \alpha \in \text{dom}(s') \setminus \delta, \exists \langle \beta, y \rangle \in X_{\langle \alpha, x \rangle}, (\langle \beta, y \rangle, [\beta', \gamma']) \in s^{**}\}.$$

We then argue that $s'' := s^{**} \cup s' \cup S_0$ is a condition by verifying the more substantial clauses from Definition 11.

Clause 2: Fix $(\langle \alpha, x \rangle, [\beta, \gamma])$ and $(\langle \alpha, x \rangle, [\beta', \gamma'])$ for which we will prove the clause. The clause follows from the fact that s^{**} and s' are conditions in \mathbb{S} if $\alpha < \delta$ or if both bits are in s' . Suppose that $(\langle \alpha, x \rangle, [\beta, \gamma]) \in s'$ and $(\langle \alpha, x \rangle, [\beta', \gamma']) \in S_0 \setminus s'$ as witnessed by $\langle \bar{\beta}, y \rangle \in X_{\langle \alpha, x \rangle}$. Since $\gamma' < \bar{\beta}$, we can assume without loss of generality that $\beta < \bar{\beta}$, from which it follows that $\gamma < \bar{\beta}$ since $s' \Vdash \text{“}\bar{\beta} \in \lim \dot{C}_\alpha^x\text{”}$. Then $(\langle \bar{\beta}, y \rangle, [\beta, \gamma]) \in s^*$ and $(\langle \bar{\beta}, y \rangle, [\beta', \gamma']) \in s^{**}$, meaning that both bits are in s^{**} . Hence it must be that either $\beta = \beta'$ or $[\beta, \gamma] \cap [\beta', \gamma'] = \emptyset$. The remaining case, in which both $(\langle \alpha, x \rangle, [\beta, \gamma])$ and $(\langle \alpha, x \rangle, [\beta', \gamma'])$ are both in $S_0 \setminus s'$, is similar.

Clause 8: Fix $\langle \alpha, x \rangle$ and β as in the statement of Clause 8 in Definition 11. If $\alpha < \delta$, then if $(\langle \alpha, x \rangle, [\beta', \gamma']) \in s^{**} \cup s' \cup S_0$ for some x, β, γ , then $(\langle \alpha, x \rangle, [\beta', \gamma']) \in s^{**}$, so we have what we need because $s^{**} \in \mathbb{S}$ and satisfies Clause 8. If $\alpha \in \aleph_2 \setminus \delta$ and $\beta \geq \delta$, then we have what we need because $s' \in \mathbb{S}$. Observe that if $\beta < \delta$, $\alpha \in \aleph_2 \setminus \delta$, and $s' \Vdash \text{“}\beta \in \lim \dot{C}_\alpha^x\text{”}$, then we have $\beta \in \text{dom}(s^{**})$, so the clause holds for the remaining case due to bits in S_0 . \square

Lemma 21. *\mathbb{S} adds a $\square(\aleph_2, \aleph_0)$ -sequence.*

Proof. Suppose G is \mathbb{S} -generic over V . We know from Lemma 17 and Corollary 20 that \mathbb{S} preserves cardinals and cofinalities up to and including \aleph_2 . In $V[G]$ we let $\vec{C} = \langle C_\alpha : \alpha \in \text{Lim}(\aleph_2) \rangle$ be defined so that $C_\alpha = \{C_\alpha^x : x \in \omega\}$ and $C_\alpha^x = \dot{C}_\alpha^x[G]$. That the C_α^x 's are clubs was handled by Proposition 13.

First, we show that \vec{C} is coherent. Suppose that $s \in \mathbb{S}$ forces that β is a limit point of \dot{C}_α^x , and suppose for contradiction that there is no $y \in \omega$ and no $s' \leq s$ forcing that $\text{“}\dot{C}_\alpha^x \cap \beta = \dot{C}_\beta^y\text{”}$. Then we find an $\leq_{\mathbb{S}}$ -decreasing sequence $\langle s_i : i < \omega \rangle$ of complete conditions such that $s_i \Vdash \text{“}\dot{C}_\alpha^x \cap \beta \neq \dot{C}_\beta^i\text{”}$,

and we find a lower bound \bar{s} of this sequence using the countable closure of \mathbb{S} . Then \bar{s} violates Clause 8.

Next, we show that $\dot{\mathcal{C}}$ does not have a thread in $V[\mathbb{S}]$, using the usual genericity argument. Suppose s forces that \dot{D} is a closed unbounded subset of \aleph_2 . Let $\bar{\alpha}$ and $s' \leq s$ be such that s' forces $\bar{\alpha}$ to be the ω^{th} point of \dot{D} . Build an \mathbb{S} -decreasing sequence of complete conditions $\langle s_i : i < \omega \rangle$ below s' and an increasing sequence of ordinals $\langle \alpha_i : i < \omega \rangle$ above $\bar{\alpha}$ as follows: Let $\alpha_0 = \bar{\alpha}$ and $s_0 = s'$. Given s_i and α_i , $s_{i+1} \leq s_i$ and α_{i+1} will be chosen such that $s_{i+1} \Vdash \text{“}\alpha_{i+1} \in \lim \dot{D}\text{”}$ and such that $\alpha_{i+1} > \max s_i$ and $\max s_{i+1} > \alpha_i$. Then let α^* be the supremum of the α_i 's and let s^* be a lower bound of the s_i 's forcing that $\dot{\mathcal{C}}_{\alpha^*}$ consists only of ω -sequences—this is specifically possible from the argument in Lemma 17, where α^* would be in $\text{fd} \setminus (\text{ed} \cup \text{fp})$ in that argument. Therefore $s^* \Vdash \text{“}\dot{D} \cap \alpha^* \notin \dot{\mathcal{C}}_{\alpha^*}\text{”}$. \square

3. THREADS

Next we introduce the threading forcing \mathbb{T} . Of course, a $\square(\aleph_2, \aleph_0)$ -sequence is defined as such because it has no thread, but our ability to force a thread allows us to make use of large cardinals from the ground model. This is also how the threading forcing for Jensen's method works. The goal of this section is to show that we can use our own \mathbb{T} to lift weakly compact embeddings of the form $j : \mathcal{M} \rightarrow \mathcal{N}$.

Definition 22. Let $G_{\mathbb{S}}$ be \mathbb{S} -generic over V and work in $V[G_{\mathbb{S}}]$. For all $\alpha \in \text{Lim}(\aleph_2)$ and $x \in \omega$, let $C_{\alpha}^x = \dot{C}_{\alpha}^x[G_{\mathbb{S}}]$.

Define \mathfrak{I} to be the set of closed intervals $[\beta, \gamma]$ such that $-1 \leq \beta \leq \gamma < \aleph_2$. Then we define a poset of \mathbb{T} of conditions $t \subset \mathfrak{I}$ such that the following holds:

- (1) The set t is countable.
- (2) If $[\beta, \gamma], [\beta', \gamma'] \in t$, then either $\beta = \beta'$ or $[\beta, \gamma] \cap [\beta', \gamma'] = \emptyset$.
- (3) If $\{[\beta_i, \gamma_i] : i < \omega\} \subseteq t$ and $\beta^* = \sup_{i < \omega} \beta_i$, then it follows that $[\beta^*, \gamma^*] \in t$ for some γ^* .
- (4) If $[\beta, \gamma] \in t$, then one of the following holds:
 - (a) $\text{cf}(\beta) = \omega_1$.
 - (b) $\beta = \beta' + 1$ and there is some $\bar{\beta} \leq \beta'$ such that $[\bar{\beta}, \beta'] \in t$.
 - (c) $\text{cf}(\beta) = \omega$ and there exists a sequence of ordinals $\langle \beta_i : i < \omega \rangle$ converging to β such that $\forall i < \omega, \exists \gamma_i, [\beta_i, \gamma_i] \in t$.
 - (d) $\text{cf}(\beta) = \omega$ and there is some $\bar{\beta} < \beta$ and some sequence of ordinals $\langle \gamma_i : i < \omega \rangle$ converging to β such that $\forall i < \omega, [\bar{\beta}, \gamma_i] \in t$.
- (5) There is some $\alpha < \aleph_2$ such that $[\alpha, \alpha] \in t$ and such that for all $\beta > \alpha$, there is no γ such that $[\beta, \gamma] \in t$. Either $\text{cf}(\alpha) = \omega_1$ or there is some $\langle \beta_n : n < \omega \rangle$ cofinal in α such that for all $n < \omega$, $[\beta_n, \gamma] \in t$ for some γ .
- (6) Suppose that $[\beta, \gamma] \in t$ and either $\text{cf}(\beta) = \omega_1$ or else $\{\beta' < \beta : \exists \gamma', [\beta', \gamma'] \in t\}$ is unbounded in β . Then there some $x \in \omega$ such that for all $\beta' \leq \gamma' < \beta$, $[\beta', \gamma'] \in t$ implies both $\beta' \in C_{\beta}^x$ and $C_{\beta}^x \cap (\beta', \gamma'] = \emptyset$.

We have $t' \leq_{\mathbb{T}} t$ if and only if:

- (1) $t' \supseteq t$;
- (2) If $\langle \alpha, x \rangle$ is such that for $\beta, \gamma < \alpha$, $[\beta, \gamma] \in t$ implies $\beta \in C_{\alpha}^x$ and $C_{\alpha}^x \cap (\beta, \gamma) = \emptyset$, then for $\beta, \gamma < \alpha$, $[\beta, \gamma] \in t'$ implies $\beta \in C_{\alpha}^x$ and $C_{\alpha}^x \cap (\beta, \gamma) = \emptyset$.

We use the following conventions for $t \in \mathbb{T}$:

- $\text{pos}(t) = \{\beta \geq 0 : \exists \gamma, [\beta, \gamma] \in t\}$.
- $\max(t)$ is the largest element of $\text{pos}(t)$.

Definition 23. If $s \in \mathbb{S}$, let

$$\text{thr}(s) = \{(\{[\alpha, \alpha]\} \cup \{[\beta, \gamma] : (\langle \alpha, x \rangle, [\beta, \gamma]) \in s\}) : \alpha \in \text{dom}(s), x \in \omega\}.$$

We say that $\langle \alpha, x \rangle$ *witnesses* that $t \in \text{thr}(s)$ if $[\alpha, \alpha] \in t$ and for $\beta, \gamma < \alpha$, $[\beta, \gamma] \in t$ if and only if $(\langle \alpha, x \rangle, [\beta, \gamma]) \in s$.

Now we can connect the notion of $\text{thr}(s)$ for $s \in \mathbb{S}$ with \mathbb{T} if we are working in V .

Proposition 24. *The following are true given $s \in \mathbb{S}$:*

- (1) *If $t \in \text{thr}(s)$ is witnessed by $\langle \alpha, x \rangle$ then $s \Vdash "t \in \dot{\mathbb{T}}"$.*
- (2) *If $s \Vdash "t \in \dot{\mathbb{T}}"$, then there is some $s' \leq s$ and some $t' \in \text{thr}(s')$ such that $s' \Vdash "t' \leq t"$.*

Proof. (1) is clear, with the appropriate $\alpha \in \text{dom } s$ and $x \in \omega$ witnessing Clause 6 from Definition 22. (2) works as follows: Let $s^* \leq s$, α , and $x \in \omega$ be such that $s^* \Vdash "\alpha = \max(\dot{t})$ and $\langle \alpha, x \rangle$ witnesses Clause 6 of Definition 22". Then use the countable closure of \mathbb{S} to find $s' \leq s^*$ such that $s' \Vdash "t = t^+ \in V"$ and such that for all $[\beta, \gamma] \in t^+$, either $(\langle \alpha, x \rangle, [\beta, \gamma]) \in s'$ or else there is some $(\langle \alpha, x \rangle, [\beta', \gamma']) \in s'$ such that $\beta \neq \beta'$ and $[\beta, \gamma] \cap [\beta', \gamma'] \neq \emptyset$. Then it must be the case that if $[\beta, \gamma] \in t^+$ then $(\langle \alpha, x \rangle, [\beta, \gamma]) \in s'$ because otherwise the definition of $\leq_{\dot{\mathbb{T}}}$ would be violated. Then let t' be such that $[\alpha, \alpha] \in t'$ and such that for all $\beta, \gamma < \alpha$, $[\beta, \gamma] \in t'$ if and only if $(\langle \alpha, x \rangle, [\beta, \gamma]) \in s'$. Then s' and t' witness (2). \square

The next step is to introduce our version of the two-step iterations that appear in Jensen's method of forcing squares.

Definition 25. We let $D(\mathbb{S} * \mathbb{T})$ refer to the set of pairs $(s, \dot{t}) \in \mathbb{S} * \mathbb{T}$ such that:

- (1) $s \Vdash "\dot{t} = \check{t}"$ for some $t \in \text{thr}(s)$.
- (2) $s \Vdash "\max(\dot{t}) = \max(s) \geq \sup(\text{out}(s))"$.

If $(s, \dot{t}) \in D(\mathbb{S} * \mathbb{T})$, we will most often write (s, \dot{t}) as (s, t) .

Lemma 26. *The set $D(\mathbb{S} * \mathbb{T})$ is dense in $\mathbb{S} * \mathbb{T}$ and countably closed. Moreover, if a sequence $\langle (s_i, t_i) : i < \omega \rangle$ is $\leq_{\mathbb{S} * \mathbb{T}}$ -decreasing, then it has a lower bound (\bar{s}, \bar{t}) such that \bar{s} is a parsimonious lower bound of $\langle s_i : i < \omega \rangle$.*

Proof. We will prove each claim separately. The "moreover" part of the statement will be clear from the proof.

Claim 27. $D(\mathbb{S} * \mathbb{T})$ is dense in $\mathbb{S} * \mathbb{T}$.

Suppose $(s, \dot{t}) \in \mathbb{S} * \mathbb{T}$. Let s' and t' witness Proposition 24 with respect to (s, \dot{t}) . Then choose $s'' \leq s'$, $t'' \in \text{thr}(s'')$, and a large enough α^* such that $(s'', \dot{t}'') \leq (s', \dot{t}')$ and such that $\alpha^* = \max(t'') = \max(s'') \geq \sup(\text{out}(s''))$ as follows: Let $\beta^* = \sup\{\gamma + 1 : \exists \beta, [\beta, \gamma] \in t'\}$ and choose α^* such that $\max(s') \cup \sup(\text{out}(s')) < \alpha^*$. Let $\langle \alpha_n : n < \omega \rangle$ be a sequence with supremum α^* such that $\alpha_0 = \beta^*$ and α_n is a successor ordinal for $n > 0$. Let s'' be

$$s' \cup \{(\langle \alpha^*, x \rangle, [\alpha_n, \alpha_{n+1} - 1]) : x, n \in \omega\} \cup \{(\langle \alpha^*, x \rangle, [\beta, \gamma]) : [\beta, \gamma] \in t', x \in \omega\}.$$

Let t'' be such that $[\alpha^*, \alpha^*] \in t''$ and such that for all $\beta, \gamma < \alpha^*$, $[\beta, \gamma] \in t''$ if and only if $(\langle \alpha^*, x \rangle, [\beta, \gamma]) \in s''$ for all $x \in \omega$.

Claim 28. $D(\mathbb{S} * \mathbb{T})$ is countably closed.

Let $\langle (s_i, t_i) : i < \omega \rangle$ be $\leq_{\mathbb{S} * \mathbb{T}}$ -decreasing in $D(\mathbb{S} * \mathbb{T})$. Let s^* be a parsimonious lower bound of $\vec{s} = \langle s_i : i < \omega \rangle$. We use some definitions with the goal of eventually constructing \bar{t} :

- $T_0 = \bigcup_{i < \omega} t_i$,
- $\text{ep}(t) = \{\beta : \exists \gamma, [\beta, \gamma] \in T_0\}$,
- $\text{fp}(t) = \lim^+(\text{ep}(t)) \setminus \text{ep}(t)$,
- $t^* = T_0 \cup \{[\beta, \beta] : \beta \in \text{fp}(t)\}$.

We will define (\bar{s}, \bar{t}) based on (s^*, t^*) and argue that $(\bar{s}, \bar{t}) \in D(\mathbb{S} * \mathbb{T})$. Now let $\bar{\alpha}$ be the supremum of $\text{ep}(t) \cup \text{fp}(t)$. If there is some i such that $\bar{\alpha} \in \text{dom}(s_i)$, in other words $\bar{\alpha} \in \text{ep}(t)$, then it will be the case that $\text{pos}(t^*) = \text{pos}(\bar{t})$ for some $\tilde{t} \in \text{thr}(\bar{s})$. Then we can let $\bar{s} = s^*$ and let \bar{t} be \tilde{t} . Let us therefore assume for the rest of the proof that for all $i < \omega$, $\bar{\alpha} \notin \text{dom } s_i$.

We will define \bar{s} as follows:

- If $\alpha \neq \bar{\alpha}$, then $(\langle \alpha, x \rangle, [\beta, \gamma]) \in s^+$ if and only if $(\langle \alpha, x \rangle, [\beta, \gamma]) \in s^*$.
- $(\langle \bar{\alpha}, x + 1 \rangle, [\beta, \gamma]) \in s^+$ if and only if $(\langle \bar{\alpha}, x \rangle, [\beta, \gamma]) \in s^*$.
- $(\langle \bar{\alpha}, 0 \rangle, [\beta, \gamma]) \in s^+$ if $[\beta, \gamma] \in t^*$ and $\gamma < \bar{\alpha}$.

We need to argue that $\bar{s} \in \mathbb{S}$. It suffices to show that Clause 8, i.e. coherence, from Definition 11 holds for bits of the form $(\langle \bar{\alpha}, 0 \rangle, [\beta', \gamma'])$. Let β be as in the statement of Clause 8. Observe that there is some $\beta^* \in (\beta, \alpha)$ such that $\beta^* \in \text{ep}(t)$ and either Case (a) or Case (c) of Clause 4 of Definition 22 holds for β^* . Let x be such that $\langle \beta^*, x \rangle$ witnesses Clause 6 of Definition 22. Then $\beta^* \in \text{dom } s^*$ and $s^* \Vdash \beta \in \lim \dot{C}_{\beta^*}^x$, so let y witness coherence of $\dot{C}_{\beta^*}^x$ at β . Then y witnesses coherence of $\dot{C}_{\bar{\alpha}}^0$ at β .

We let \bar{t} be defined so that $[\bar{\alpha}, \bar{\alpha}] \in \bar{t}$ and so that for all $\beta, \gamma < \bar{\alpha}$, $[\beta, \gamma] \in \bar{t}$ if and only if $(\langle \bar{\alpha}, 0 \rangle, [\beta, \gamma]) \in \bar{s}$. We claim that $(\bar{s}, \bar{t}) \in D(\mathbb{S} * \mathbb{T})$. It can be checked that \bar{s} is a lower bound of $\langle s_i : i < \omega \rangle$: Observe that $\bar{\alpha} = \max(\bar{s}) = \max(s^*)$ because s^* was chosen to be parsimonious and because Clause 2 and Clause 1 from Definition 25 hold for (s_i, t_i) for all $i < \omega$. In particular, the fact that $\sup(\text{out}(s_i)) \leq \max(s_i)$ for all $i < \omega$ implies that $\text{fp}(\bar{s}) \subseteq \max(s^*)$. We only needed to guarantee that we would have $\bar{t} \in \text{thr}(\bar{s})$. \square

We therefore see that \mathbb{T} preserves \aleph_1 over $V[\mathbb{S}]$ since it is a factor of a countably distributive iteration. Moreover, we can expand the proof from Lemma 19 to see:

Proposition 29. $\mathbb{S} * \mathbb{T}$ preserves \aleph_2 over V .

Proof. Specifically, suppose that $s \in \mathbb{S}$, $t \in \mathbb{T}$, and $M \prec K$ is a basic model with $s, \mathbb{S} * \mathbb{T} \in M$, and $\delta = M \cap \aleph_2$. Then it can be argued that $(s, t \cup [\delta, \delta])$ is strongly $(M, \mathbb{S} * \mathbb{T})$ -generic. Then apply Proposition 6. \square

Although \mathbb{T} preserves cardinals, it does not preserve the canonical $\square(\aleph_2, \aleph_0)$ -sequence added by \mathbb{S} . This is the primary function of \mathbb{T} in some sense, despite its usefulness for lifting embeddings (which will be established shortly).

Proposition 30. If $G_{\mathbb{S}} * G_{\mathbb{T}}$ is $\mathbb{S} * \mathbb{T}$ -generic over V , and $\mathcal{D} = \{\beta < \aleph_2 : \exists \gamma, [\beta, \gamma] \in G_{\mathbb{T}}\}$, then \mathcal{D} is a thread of the $\square(\aleph_2, \aleph_0)$ -sequence \mathcal{C} derived from $G_{\mathbb{S}}$.

The following lemma shows how we will use a weakly compact cardinal together with \mathbb{S} by lifting embeddings, using \mathbb{T} in a crucial way. In the context of Jensen's method for forcing squares, we would use a generic condition argument for $j(\mathbb{S})$. However, when we have lifted $j : \mathcal{M}[\text{Col}(\aleph_1, < \kappa)] \rightarrow \mathcal{N}[\text{Col}(\aleph_1, < j(\kappa))]$ and are working in $V[\text{Col}(\aleph_1, < j(\kappa))]$, we have $|\mathbb{S}| = \aleph_1$, yet the conditions in both \mathbb{S} and $j(\mathbb{S})$ are countable. Therefore we use a factorization argument instead.

Lemma 31. Suppose the following:

- $\mathfrak{C} = \text{Col}(\aleph_1, < \kappa)$ is the Lévy collapse and $G_{\mathfrak{C}}$ is \mathfrak{C} -generic over \bar{V} .
- $j : \mathcal{M} \rightarrow \mathcal{N}$ is a weakly compact embedding with critical point κ .
- $j : \mathcal{M}[G_{\mathfrak{C}}] \rightarrow \mathcal{N}[j(G_{\mathfrak{C}})]$ is the usual lifted embedding given by $\dot{x}_{G_{\mathfrak{C}}} \mapsto j(\dot{x})_{j(G_{\mathfrak{C}})}$.
- $V[j(G_{\mathfrak{C}})]$ contains generics $G_{\mathbb{S}}$ and $G_{\mathbb{T}}$ for \mathbb{S} and \mathbb{T} respectively.

Then in $\bar{V}[j(G_{\mathfrak{C}})]$, $j(\mathbb{S})$ is forcing equivalent to $\mathbb{S} * \mathbb{T} * \mathbb{S}'$ where \mathbb{S}' is a countably closed forcing over $\bar{V}[j(G_{\mathfrak{C}})]$.

Observe that the fourth premise can be fulfilled using the Absorption Lemma.

Proof. First we describe \mathbb{S}' and some relevant dense subsets of \mathbb{S} and $j(\mathbb{S})$. Working in $V[G_{j(\mathfrak{C})}]$, $j(\mathfrak{S})$ (where \mathfrak{S} is from Definition 11) is of course equal to the set of bits $(\langle \alpha, x \rangle, [\beta, \gamma])$ such that $\alpha < j(\kappa)$, $x \in \omega$, $\beta \leq \gamma < j(\kappa)$. Let $C_{\alpha}^x = \dot{C}_{\alpha}^x[G_{\mathbb{S}}]$ for all $\langle \alpha, x \rangle \in \text{Lim}(\aleph_2) \times \omega$ and let $\mathcal{D} = \{\beta < \aleph_2 : \exists \gamma, [\beta, \gamma] \in G_{\mathbb{T}}\}$.

Now let \mathbb{S}' consist of $s \subseteq j(\mathfrak{S}) \setminus \mathfrak{S}$ such that Clause 1 through Clause 7 from Definition 11 hold, such that its ordering is defined like $\leq_{\mathbb{S}}$, and moreover two additional clauses hold, the first of which is a modification of Clause 8 from Definition 11:

- (10) Suppose that $(\langle \alpha, x \rangle, [\beta, \gamma]) \in s$ and either $\text{cf}(\beta) = \omega_1$ or else $\{\beta' < \beta : \exists \gamma', (\langle \alpha, x \rangle, [\beta', \gamma']) \in s\}$ is unbounded in β . Then:
 - (a) If $\beta > \kappa$, then there is some $y \in \omega$ such that for all $\beta' \leq \gamma' < \beta$, $(\langle \alpha, x \rangle, [\beta', \gamma']) \in s$ if and only if $(\langle \beta, y \rangle, [\beta', \gamma']) \in s$.

- (b) If $\beta < \kappa$, then there is some $y \in \omega$ such that for all $\beta' \leq \gamma' < \beta$, if $(\langle \alpha, x \rangle, [\beta', \gamma']) \in s$ then $\beta' \in C_\beta^y$ and $C_\beta^y \cap (\beta', \gamma') = \emptyset$.
 - (c) If $\beta = \kappa$, then for all $\beta', \gamma' < \beta$, if $(\langle \alpha, x \rangle, [\beta', \gamma']) \in s$, then $\beta' \in \mathcal{D}$ and $\mathcal{D} \cap (\beta', \gamma') = \emptyset$.
- (11) For all limit ordinals α and all $x \in \omega$ such that $\exists \tilde{\beta}, \tilde{\gamma}, (\langle \alpha, x \rangle, [\tilde{\beta}, \tilde{\gamma}]) \in s$, one of the following holds:
- (a) $(\langle \alpha, x \rangle, [\kappa, \gamma]) \in s$ for some γ .
 - (b) There exists a non-empty finite sequence $\beta_0 < \beta_1 < \dots < \beta_k$ such that the following hold:
 - (i) For all $\ell \in \{0, 1, \dots, k\}$, $(\langle \alpha, x \rangle, [\beta_\ell, \gamma]) \in s$ for some γ ;
 - (ii) Either $\text{cf}(\beta_0) = \omega_1$ or there are sequences $\langle \beta_n : n < \omega \rangle$, $\langle \gamma_n : n < \omega \rangle$ with $\sup_{n < \omega} \beta_n = \beta$ and $(\langle \alpha, x \rangle, [\beta_n, \gamma_n]) \in s$ for all $n < \omega$;
 - (iii) For all β_ℓ with $\ell > 0$, either $\beta_\ell = \beta_{\ell-1} + 1$, or $\beta_\ell = \bar{\beta} + 1$ and $(\langle \alpha, x \rangle, [\beta_{\ell-1}, \bar{\beta}]) \in s$, or there is a sequence $\langle \gamma_n : n < \omega \rangle$ with $\sup_{n < \omega} \gamma_n = \beta_\ell$ and $(\langle \alpha, x \rangle, [\beta_{\ell-1}, \gamma_n]) \in s$ for all $n < \omega$.
 - (iv) $(\langle \alpha, x \rangle, [\beta_\ell, \gamma]) \in s$ for some $\gamma \geq \kappa$.

Let $D(j(\mathbb{S}))$ be the dense subset of $s \in j(\mathbb{S})$ such that Clause 11 holds for s . Now we give the isomorphism $\Psi : \{(s, t, s') : s \in \mathbb{S}, (s, t) \in D(\mathbb{S} * \mathbb{T}), s' \in \mathbb{S}'\} \rightarrow D(j(\mathbb{S}))$. Specifically,

$$\Psi(s, t, s') = s \cup s' \cup \{(\langle \kappa, x \rangle, [\beta, \gamma]) : x \in \omega, [\beta, \gamma] \in t\}.$$

Observe that the definition of Ψ is where we use Clause 3 from Definition 11. We are also using the fact that $V[G_{j(\mathfrak{e})}] \models “\kappa \in j(\kappa) \cap \text{cof}(\omega_1)”$. It is immediate that Ψ is order-preserving. Moreover, Ψ is a bijection because it has a natural inverse.

Now we will show that \mathbb{S}' is countably closed in $V[G_{j(\mathfrak{e})}]$. Suppose $\vec{s} = \langle s_i : i < \omega \rangle$ is a descending sequence in \mathbb{S}' . Define ed , fd , ep , fp as in Definition 16, and let \bar{s} be a parsimonious lower bound of \vec{s} defined exactly as in Lemma 17. So $\bar{s} \in j(\mathbb{S})$. We know that \bar{s} satisfies Clause 1 through Clause 7 from Definition 11. We will prove Clause 10 after Clause 11, since the former depends on the proof of the latter.

Clause 11: The case where $\alpha \in \text{fd} \setminus (\text{ep} \cup \text{fp})$ is trivial. Given $\alpha \in \text{ed}$, $x \in \omega$, there is some i such that $\alpha \in \text{dom}(s_i)$. Then the fact that s_i satisfies Clause 11 is enough to imply that \bar{s} satisfies Clause 11. If $\alpha \in \text{fp}$, then there is some $\langle \alpha', x' \rangle$ witnessing $\alpha \in \text{fp}$, and we get Clause 11 from the s_i for i large enough that $\alpha' \in \text{dom}(s_i)$.

Clause 10: For the purpose of verifying this clause, when we say that y witnesses coherence of C_α^x at β for $\beta \leq \kappa$, we are referring to the y mentioned in Case (b) or Case (c) of the clause.

Now fix $\alpha \in \text{Lim}(j(\kappa) \setminus (\kappa + 1))$ and $x \in \omega$ as in the statement of Clause 10. We have already handled the case where $\beta > \kappa$ in Lemma 17. Therefore we can assume that $\beta < \kappa$ for the rest of the proof. We break the proof into cases.

- $\alpha \in \mathbf{ep}, \beta \in \mathbf{ep}$, for large i **Case (a) of Clause 11 holds for s_i w.r.t. $\langle \alpha, x \rangle$** : We have that $\beta \in \lim \mathcal{D}$, so there is some $y \in \omega$ such that $\mathcal{D} \cap \beta = \dot{C}_\beta^y$. Then y witnesses coherence of \dot{C}_x^α at β .
- $\alpha \in \mathbf{ep}, \beta \in \mathbf{fp}$, for large i **Case (a) of Clause 11 holds for s_i w.r.t. $\langle \alpha, x \rangle$** : Observe that since $\mathbb{S} * \mathbb{T}$ has a countably closed dense subset, it follows that \mathbb{T} is countably distributive, so $G_{\mathbb{T}}$ is countably closed, hence $\beta \in \lim \mathcal{D}$. This then follows the reasoning of the previous case.
- $\alpha \in \mathbf{ep}, \beta \in \mathbf{ep}$, for large i **Case (b) of Clause 11 holds for s_i w.r.t. $\langle \alpha, x \rangle$** : This means that there is some β^+ such that for large i , $s_i \Vdash \text{“}\beta^+ = \max(\lim \dot{C}_\alpha^x \cap \kappa)\text{”}$, and that $\beta \leq \beta^+$. If $\beta = \beta^+$ and y witnesses coherence of \dot{C}_α^x at β^+ , then we are done with this case. If $\beta < \beta^+$, y witnesses coherence of \dot{C}_α^x at β^+ , and y' is such that $C_{\beta^+}^y \cap \beta = C_\beta^{y'}$ (since $\beta \in \lim C_{\beta^+}^y$), then y' witnesses coherence of \dot{C}_α^x at β .
- $\alpha \in \mathbf{ep}, \beta \in \mathbf{fp}$, for large i **Case (b) of Clause 11 holds for s_i w.r.t. $\langle \alpha, x \rangle$** : This is like the previous case, noting that if $\beta < \beta^+$ as written there, then we would still have $\beta \in \lim C_{\beta^+}^y$.
- $\langle \tilde{\alpha}, \tilde{x} \rangle$ witnesses $\alpha \in \mathbf{fp}, \beta \in \mathbf{ep}$, for lg. i **Case (a) of Clause 11 holds for s_i , $\langle \tilde{\alpha}, \tilde{x} \rangle$** : We have $\beta \in \lim \mathcal{D}$, so there is some $y \in \omega$ such that $\mathcal{D} \cap \beta = \dot{C}_\beta^y$. So y witnesses coherence of $\dot{C}_{\tilde{\alpha}}^{\tilde{x}}$ at β , hence it witnesses coherence of C_α^x at β .

The remaining cases all consider $\alpha \in \mathbf{fp}$ and take $\langle \tilde{\alpha}, \tilde{x} \rangle$ witnessing this. The reasoning depends on whether Case (a) or Case (b) holds for s_i with respect to $\langle \tilde{\alpha}, \tilde{x} \rangle$ for large i . The arguments are analogous to the cases already discussed. \square

Now we can use \mathbb{T} to show that stronger squares do not hold. Since the bulk of the work was done with Lemma 31, and the remaining lemmas are variations of standard arguments, we will handle the rest with a light amount of detail.

Lemma 32. *If W is a ground model and \mathcal{C} is a $\square(\aleph_2, < \aleph_0)$ -sequence in $W[\mathbb{S}]$, then \mathbb{T} does not thread \mathcal{C} .*

Proof. This lemma has analogs for Jensen-style posets (see [8], Lemma 4.5). Let \dot{D} be a name for a thread for \mathcal{C} added over $W[\mathbb{S}]$ by \mathbb{T} and work in W . Build a descending sequence $\langle s_i : i < \omega \rangle$ of conditions in \mathbb{S} , sequences $\langle t_i^j : j < \omega \rangle$ in \mathbb{T} for $i < \omega$, and an increasing sequence of ordinals $\langle \gamma_i : i < \omega \rangle$ such that the following hold:

- $(s_i, t_i^j) \in D(\mathbb{S} * \mathbb{T})$ for all $i, j < \omega$.
- $t_i^j = t_0^j$ for all $j < i$.
- For all $i, j < \omega$, $\max t_0^j = \max t_i^j$.
- (s_{i+1}, t_0^{i+1}) and (s_{i+1}, t_{i+1}^{i+1}) decide “ $\gamma_i \in \dot{D}$ ” differently.

Then let \bar{s} be a lower bound of $\langle s_i : i < \omega \rangle$ and let \bar{t}_i be a lower bound of $\langle t_i^j : j < \omega \rangle$ for all $i < \omega$, and also let $\bar{\gamma}$ be a lower bound of $\langle \gamma_i : i < \omega \rangle$. Then each (\bar{s}, \bar{t}_i) gives a different possibility for $C \in \mathcal{C}_{\bar{\gamma}}$, contradicting the fact that \mathcal{C} is a $\square(\aleph_2, < \aleph_0)$ -sequence. \square

Lemma 33. *If \mathcal{C} is a $\square(\lambda, \aleph_0)$ -sequence and \mathbb{P} is countably closed, then \mathbb{P} does not thread \mathcal{C} .*

Proof. Stronger versions of this lemma appear elsewhere ([6]), but we sketch an argument for completeness.

Let \dot{D} be a name for a thread supposedly added by \mathbb{P} . Build a tree $\langle p_x : x \in 2^{<\omega} \rangle$ and a collection of ordinals $\langle \gamma_x : x \in 2^{<\omega} \rangle$ such that:

- $x \sqsubseteq y$ implies $p_y \leq p_x$,
- $p_{x \smallfrown 0}$ and $p_{x \smallfrown 1}$ decide “ $\gamma_x \in \dot{D}$ ” differently for some γ_x above $\sup_{|y| < x} \gamma_y$.

Then for all $X \in 2^\omega$, let p_X be the lower bound of $\langle p_{X \upharpoonright n} : n < \omega \rangle$. Let $\gamma = \sup_{x \in 2^\omega} \gamma_x$. Then the p_X 's give 2^ω -many possibilities for $C \in \mathcal{C}_\gamma$, which is a contradiction of the fact that \mathcal{C} is a $\square(\lambda, \aleph_0)$ -sequence. \square

Proposition 34. *If \mathcal{C} is a $\square_{\mu, \kappa}$ -sequence and \mathbb{P} preserves μ^+ , then \mathbb{P} does not thread \mathcal{C} .*

Proof. This is basically a restatement of Remark 4. \square

Corollary 35. *If κ is weakly compact, then $V[\text{Col}(\aleph_1, < \kappa)][\mathbb{S}] \models \neg \square(\aleph_2, < \aleph_0)$.*

Proof. Let M be a κ -sized transitive model containing \mathbb{S} and a supposed $\text{Col}(\aleph_1, < \kappa) * \mathbb{S}$ -name \dot{C} for a $\square(\aleph_2, < \aleph_0)$ -sequence. Let $j : M \rightarrow N$ be a weakly compact embedding with critical point κ , and let $G_{\mathcal{C}} * G_{\mathbb{S}} * G_{\mathbb{T}}$ be $\text{Col}(\aleph_1, < \kappa) * \mathbb{S} * \mathbb{T}$ -generic over V . Since $\mathbb{S} * \mathbb{T}$ is countably closed, the quotient $\mathfrak{R} := \text{Col}(\aleph_1, < j(\kappa)) / G_{\mathcal{C}} * G_{\mathbb{S}} * G_{\mathbb{T}}$, is forcing-equivalent to a countably closed forcing (see the section on absorption in [3]). Let $G_{\mathfrak{R}}$ be generic for \mathfrak{R} and work in $V[G_{\mathcal{C}} * G_{\mathbb{S}} * G_{\mathbb{T}} * G_{\mathfrak{R}}]$ to use Lemma 31 to lift j to $j : M[G_{\mathcal{C}}][G_{\mathbb{S}}] \rightarrow N[j(G_{\mathcal{C}})][j(G_{\mathbb{S}})]$ by forcing with a \mathbb{S}' -generic $G_{\mathbb{S}'}$. Use the lift to define a thread D for $\mathcal{C} := \dot{C}[G_{\mathcal{C}} * G_{\mathbb{S}}]$ by taking an element from the κ^{th} level of $j(\mathcal{C})$. Lemma 32 shows that \mathbb{T} could not have added D , and Lemma 33 show that neither \mathfrak{R} nor \mathbb{S}' could have added D . Therefore the thread D already exists in $V[G_{\mathcal{C}} * G_{\mathbb{S}}]$, and so \mathcal{C} is not actually a $\square(\aleph_2, < \aleph_0)$ -sequence. \square

Corollary 36. *If κ is weakly compact then $V[\text{Col}(\aleph_1, < \kappa)][\mathbb{S}] \models \neg \square_{\aleph_1, \aleph_0}$.*

Proof. A Mahlo cardinal would be sufficient (see [8], Theorem 4.4). But assume we have a weakly compact cardinal and define a thread D as in Corollary 35. Then Proposition 34 and Proposition 29 together imply that \mathbb{T} could not have added D . Then Lemma 33 shows that the rest of the extension could not have added D . \square

Hence we have finished proving Theorem 1.

Acknowledgement. I would like to thank Heike Mildenerger for a great deal of support while I was working on this paper. I am especially grateful for her willingness to spend a lot of time looking at rough drafts. I would also like to thank the anonymous referee for carefully reading the manuscript and finding many problems in the submitted version.

REFERENCES

- [1] Uri Abraham and Saharon Shelah. Forcing closed unbounded sets. *J. Symbolic Logic*, 48(3):643–657, 1983.
- [2] James Cummings. Notes on singular cardinal combinatorics. *Notre Dame J. Formal Logic*, 46(3), 2005.
- [3] James Cummings. Iterated forcing and elementary embeddings. In Matthew Foreman and Akihiro Kanamori, editors, *Handbook of Set Theory*, pages 775–883. Springer, 2010.
- [4] Gregor Dolinar and Mirna Džamonja. Forcing \square_{ω_1} with finite conditions. *Ann. Pure Appl. Logic*, 164(1):49–64, 2013.
- [5] Todd Eisworth. Successors of singular cardinals. In Matthew Foreman and Akihiro Kanamori, editors, *Handbook of Set Theory*, pages 1229–1350. Springer, 2010.
- [6] Yair Hayut and Chris Lambie-Hanson. Simultaneous stationary reflection and square sequences. *J. Math. Log.*, 17(2):1750010, 2017.
- [7] Thomas Jech. *Set Theory*. Springer Monographs in Mathematics. Springer-Verlag, Berlin, the third millennium, revised and expanded edition, 2003.
- [8] Menachem Magidor and Chris Lambie-Hanson. On the strengths and weaknesses of weak squares. In *Appalachian set theory 2006–2012*, volume 406 of *London Math. Soc. Lecture Note Ser.*, pages 301–330. Cambridge Univ. Press, Cambridge, 2013.
- [9] William Mitchell. Adding closed unbounded subsets of ω_2 with finite forcing. *Notre Dame J. Formal Logic*, 46(3):357–371, 2005.
- [10] William J. Mitchell. On the Hamkins approximation property. *Ann. Pure Appl. Logic*, 144:126–129, 2006.
- [11] Itay Neeman. Two applications of finite side conditions at ω_2 . *Arch. Math. Logic*, 56(7-8):983–1036, 2017.
- [12] Saharon Shelah and Lee Stanley. S -forcing. I. A “black-box” theorem for morasses, with applications to super-Souslin trees. *Israel J. Math.*, 43(3):185–224, 1982.
- [13] Stevo Todorčević. Partitioning pairs of countable ordinals. *Acta Math.*, 159(3-4):261–294, 1987.

ALBERT-LUDWIGS-UNIVERSITÄT FREIBURG, MATHEMATISCHES INSTITUT, ABTEILUNG
 FÜR MATH. LOGIK, ERNST-ZERMELO-STRASSE 1, 79104 FREIBURG IM BREISGAU,
 GERMANY

Email address: maxwell.levine@mathematik.uni-freiburg.de